American Heart Journal

The international publication for the study of the heart and circulation

Editor in Chief: DEAN T. MASON, M.D.

Associate Editor: ANTHONY N. DEMARIA, M.D.

Assistant Editors: EZRA A. AMSTERDAM, M.D.

NAJAM A. AWAN, M.D. GARRETT LEE, M.D.

THOMAS A. RIEMENSCHNEIDER, M.D.

ZAKAUDDIN VERA, M.D.

Managing Editor: LESLIE J. SILVERNAIL

Editorial Board

W. Gerald Austen, M.D., Boston Daniel S. Berman, M.D., Los Angeles J. Thomas Bigger, M.D., New York William J. Bommer, M.D., Davis Lawrence I. Bonchek, M.D., Milwaukee Lawrence S. Cohen, M.D., New Haven Lawrence H. Cohn, M.D., Boston John J. Collins, Jr., M.D., Boston C. Richard Conti, M.D., Gainesville Mary Allen Engle, M.D., New York Harvey Feigenbaum, M.D., Indianapolis James M. Foerster, M.D., Davis Victor F. Froelicher, M.D., San Diego Paul C. Gillette, M.D., Houston John F. Goodwin, M.D., London Antonio M. Gotto, Jr., M.D., Houston Rolf M. Gunnar, M.D., Maywood Fred J. Harris, M.D., Davis W. Proctor Harvey, M.D., Washington, D.C. Richard H. Helfant, M.D., Philadelphia Walter L. Henry, M.D., Irvine J. O'Neal Humphries, Columbia J. Willis Hurst, M.D., Atlanta Mark E. Josephson, M.D., Philadelphia James A. Joye, M.D., Davis

William B. Kannel, M.D., Framingham Norman M. Kaplan, M.D., Dallas Samuel Kaplan, M.D., Cincinnati Joseph A. Kisslo, M.D., Durham Lawrence J. Laslett, M.D. Davis Michael Lesch, M.D., Chicago Floyd D. Loop, M.D., Cleveland Reginald I. Low, M.D., Davis Robert I. Levy, M.D., Bethesda Dan G. McNamara, M.D., Houston Richard R. Miller, M.D., Houston Robert A. O'Rourke, M.D., San Antonio Joseph K. Perloff, M.D., Los Angeles Bertram Pitt, M.D., Ann Arbor Charles E. Rackley, M.D., Birmingham William C. Roberts, M.D., Bethesda David J. Sahn, M.D., Tucson Robert C. Schlant, M.D., Atlanta Pravin M. Shah, M.D., Los Angeles Walter Sommerville, M.D., London James T. Willerson, M.D., Dallas David O. Williams, M.D., Providence Roger A. Winkle, M.D., Stanford Robert F. Zelis, M.D., Hershey Douglas P. Zipes, M.D., Indianapolis

VOLUME 103

JANUARY-JUNE 1982

VOLUME 103

COPYRIGHT® 1982 BY

THE C.V. MOSBY COMPANY

All rights reserved

Printed in the United States of America

Author index*

A

ABRAHAM, ABRAHAM K., AND NORRIS, ROBIN M. Sequential AV pacing in right ventricular myocardial infarction with complete heart block, 929 (Letter to Editor)

ACCETTO, BOJAN. Beneficial hemorheologic therapy of chronic peripheral arterial disorders with pentoxifylline: Results of double-blind study versus vasodilator-nyli-

ACHUFF, STEPHEN C. (See Heuser, Achuff, and Brinker), 430
ALEXANDER, JONATHAN. (See Lewis et al.), 498
ALLEN, HUGH D. (See Valdes-Cruz et al.), 505
— (See Goldberg et al.), 858
AMSTERDAM, EZRA A. (See Dressendorfer et al.), 358

AMSTERDAM, EZRA A. (See Dressendorfer et al.), 358

—. (See Awan et al.), 474

Anderson, Charles M. (See Bragg-Remschel, Anderson, and Winkle), 20

Andrews, Virginia. (See Weber, Andrews, and Janicki), 639

Antonelli, R. (See Biagini et al.), 13

Ariet, Mario. (See Geiser et al.), 1056

ARONOW, WILBERT S. (See Butman et al.), 912
ARTMAN, MICHAEL, AND GRAHAM, THOMAS P., JR. Congestive heart failure in infancy: Recognition and management,

ARYANPUR-KASHANI, IRAJ. Verapamil effects on retrograde conduction in supraventricular tachycardia therapy, 306 (Letter to Editor)

Awan, Najam A. (See Vera, Awan, and Mason), 80

—, Amsterdam, Ezra A., Hermanovich, John, Bommer, William
J., Needham, Kathleen E., and Mason, Dean T.
Long-term hemodynamic and clinical efficacy of captopril therapy in ambulatory management of severe chronic congestive heart failure, 474

AYRES, STEPHEN M. Mechanisms and consequences of pulmonary edema: Cardiac lung, shock lung, and principles of ventilatory therapy in adult respiratory distress syn-

Badke, Frederick R. (See O'Rourke, Badke, and Forst), 489 Bailey, John C. (See Lindemann, Bailey, and Watanabe), 746 Baker, John T. (See Bardy et al.), 1068 Baker, Mark. (See Moncada et al.), 263 Baky, Sherrin. (See Nademanee et al.), 802 Bar, Frits W. H. M. (See de Zwaan, Bär, and Wellens), 730

drome, 97

Baran, Andril, Nanda, Navin C., Falkorf, Michael, Barold, S. Serge, and Gallaguer, John J. Two-dimensional echocardiographic detection of arrhythmogenic right ventricular dysplasia, 1066 (Brief comm.)

BARDY, GUST H., VALENSTEIN, PAUL, STACK, RICHARD S., BAKER, JOHN T., AND KISSLO, JOSEPH A. Two-dimensional echocardiographic identification of sinus of Valsalvaright heart fistula due to infective endocarditis, 1068 (Brief comm.)

(Brief comm.)

BARLOW, JOHN B. (See Davis et al.), 289

BAROLD, S. SERGE. (See Ong, Nanda, and Barold), 917

—. (See Baran et al.), 1066

BARRIO, JORGE. (See Schön et al.), 532

BARRIO, JORGE. (See Schion et al.), 502

—. (See Schön et al.), 548

BARRY, WILLIAM H. (See Smith et al.), 716

BATEMAN, TIM. (See Schiff et al.), 562

BAUERNPEIND, ROBERT A. (See Swiryn et al.), 168

UERNFEIND, ROBERT A. (See Swiftyr et al.), 400 (See Swiryn et al.), 319 (See Strasberg et al.), 795 Swiryn, Steven P., Strasberg, Boris, Palileo, Edwin, Sca-Gliotti, Daniel, and Rosen, Kenneth M. Electrophysiologic drug testing in prophylaxis of sporadic paroxysmal atrial fibrillation: Technique, application, and efficacy in patients, 941 severely symptomatic preexcitation

*January, pp. 1-160; February, pp. 161-318; March, pp. 319-458; April, Part 1, pp. 459-582; April, Part 2, pp. 583-788; May, pp. 789-940; June, pp. 941-1128

BEDER, STANLEY D., NIHILL, MICHAEL R., AND MCNAMARA, DAN G. Idiopathic dilation of the right atrium in a child, 134 (Brief comm.)

-, GILLETTE, PAUL C., GARSON, ARTHUR, JR., AND MCNAMARA, DAN G. Clinical confirmation of ECG criteria for left atrial rhythm, 848

BELL, WILLIAM R., AND SIMON, TOBY L. Current status of pulmonary thromboembolic disease: Pathophysiology, diagnosis, prevention, and treatment, 239

Ben-Ishay, D. (See Yagil et al.), 310

Bennett, David H. (See Ramsdale et al.), 459

(See Ramsdale and Bennett), 1083

BERESFORD-KROEGER, D. (See Biro, Beresford-Kroeger, and Hendry), 870

BERGER, HARVEY J. (See Lewis et al.), 498 BERMAN, DANIEL. (See Schiff et al.), 562

BERNSTEIN, KURT. (See Ulmsten, Bernstein, and Mattson), 448

DERINSTEIN, KUIFT. (See Utimstein, Derinstein, and Mattson), 448
BERTORINI, TULIO F. (See Gottdiener et al.), 525
BESLEY, DENNIS C., McWilliams, Gregory J., Moodie, Douglas S.,
AND CASTLE, LON W. Long-term follow-up of young
adults following permanent pacemaker placement for
complete heart block, 332

Besozzi, Myrwood C. (See Rizi et al.), 292
Besterman, Edwin. Waller before Einthoven in the clinical development of noninvasive ECG, 572 (Letter to Editor)

BETHGE, CHRISTIAN. (See Meyer et al.), 132

BETHGE, CHRISTIAN. (See Meyer et al.), 132

BIAGINI, A., MAZZEI, M. G., CARFEGGIANI, C., TESTA, R., ANTONELLI, R., MICHELASSI, C., L'ABBATE, A., AND MASERI, A. Vasospastic ischemic mechanism of frequent asymptomatic transient ST-T changes during continuous electrocardiographic monitoring in selected unstable

electrocardiographic includes a grant angina patients, 13

Biancaniello, Thomas M., Meyer, Richard A., Gaum, Winston E., and Kaplan, Samuel. Primary benign intramural ventricular tumors in children: Pre- and postoperative electrocardiographic, echocardiographic and angiocar-diographic evaluation, 852

BIGGER, J. THOMAS, JR. (See Goldberg et al.), 75

—, WELD, FRANCIS M., AND ROLNITZKY, LINDA M. Which postinfarction ventricular arrhythmias should be treated,

BIRO, GEORGE P., BERESFORD-KROEGER, D., AND HENDRY, P. Early deleterious hemorheologic changes following acute experimental coronary occlusion and salutary antihyperviscosity effect of hemodilution with stroma-free
hemoglobin, 870

BISHOP, RICHARD L. (See Doi et al.), 301

BLACKMAN, MARIE S. (See Kavey, Blackman, and Sondheimer),
242

BLAKE, JEFFREY. (See Kay, Blake, and Rubin), 308 Blank, David. (See Nanji and Blank), 1088 Bommer, William J. (See Awan et al.), 474

Borow, Kennerh M., and Newburger, Jane W. Noninvasive estimation of central aortic pressure using the oscillometric method for analyzing systemic artery pulsatile blood flow: Comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure measurements, 879

BOUCHER, CHARLES A. (See Brown et al.), 1019

Bragg-Remschel, Deborah A., Anderson, Charles M., and Win-KLE, ROGER A. Frequency response characteristics of ambulatory ECG monitoring systems and their impli-cations for ST segment analysis, 20

Braunwald, Eugene. (See Rude et al.), 986 Bray, Colin L. (See Ramsdale et al.), 459 Bream, Peter R. (See Rogers et al.), 185

Brenner, Robert A. (See Riemenschneider et al.), 834 Brinker, Jeffrey A. (See Heuser, Achuff, and Brinker), 430 Bristow, Michael R. (See Ginsburg and Bristow), 928

BROOKS, HAROLD L. (See Ptacin et al.), 54 Brown, B. Greg. (See Josephson et al.), 1008 Brown, Jerry M. (See Sobol et al.), 563 Brown, Kenneth A., Boucher, Charles A., Okada, Robert D., Strauss, H. William, and Pohost, Gerald. Initial and delayed right ventricular thallium-201 rest-imaging following dipyridamole-induced coronary vasodila-tion: Relationship to right coronary artery pathoana-

Brown, Mark. (See Covitz et al.), 1072
Browne, Keven F. (See Zipes et al.), 789
Brundage, Bruce H., and Lipton, Martin J. The emergence of computed tomography as a cardiovascular diagnostic

Computed Comography as a Cartic technique, 313

BRUNDLER, H. (See Pfisterer et al.), 92

BRUNNER, HANS R. (See Waeber et al.), 384

BUJA, L. MAXIMILJAN. (See Nicod et al.), 822

BUNKER, STEPHEN R. (See Sobol et al.), 563

Buttu, Ioan. (See Georgescu et al.), 572

Buttman, Samuel, Olson, Harold G., Aronow, Wilbert S., and
Lyons, Kennetth P. Remote right ventricular myocardial infarction mimicking chronic pericardial constriction, 912 (Brief comm.)

Byrom, Ernest. (See Swiryn et al.), 319

CALDWELL, RANDALL L. (See Hurwitz et al.), 421 CAMERON, E. A. (See Singh and Cameron), 449 CANTWELL, JOHN D. Life-threatening exertional ventricular tachy-cardia in ischemic heart disease best treated by coronary bypass surgery, 152 (Letter to Editor) Cardoso, Manuel. (See Moncada et al.), 263

CARDOSO, MANUEL. (See Moncada et al.), 263
CAREY, JOAN R. (See Johnson et al.), 4
CARPEGGIANI, C. (See Biagini et al.), 13
CASPI, AVI. (See Davis et al.), 289
CASTELLANOS, AGUSTIN. (See Myerburg et al.), 615
CASTLE, LON W. (See Besley et al.), 332
CHAIGNON, MICHEL, CHEN, WEI-TZUOH, TARAZI, ROBERT C., NAKAMOTO, SATORU, AND SALCEDO, ERNESTO. Acute effects of hemodialysis on echographic-determined cardiac performance: Improved contractility resulting from formance: Improved contractility resulting from serum increased calcium with reduced potassium

despite hypovolemic-reduced cardiac output, 374
Chan, William. (See Rizi et al.), 292
Charocopos, Fotis. (See Waeber et al.), 384
Charuzi, Yzhar. (See Valovis and Charuzi), 432
Chatterjee, Kanu. (See Parmley, Rouleau, and Chatterjee),

CHEN, WEI-TZUGH. (See Chaignon et al.), 374
CHERIAN, GEORGE, TEI, CHUWA, SHAH, PRAVIN M., AND WONG,
MAYLENE. Diastolic prolapse in the flail mitral valve
syndrome: A new observation providing differentiation from the mitral valve prolapse syndrome, 1074 (Brief comm.)

CHILD, JOHN S., AND PERLOFF, JOSEPH K. Reply, 1087 (Letter to

Editor)
CHILSON, DONALD A. (See Zipes et al.), 789
CHIPPERFIELD, BARBARA. Water hardness, myocardial metal concentrations, and sudden coronary death, 1085 (Letter

CHIRIKOS, THOMAS N. (See Lewis et al.), 123 CHRISTIE, LEONARD G., Jr. (See Geiser et al.), 1056 CHURCHILL, ROBERT. (See Moncada et al.), 263 CLARK, LELAND C., Jr. (See Rude et al.), 986 CLARK, T. J. H. (See Rees and Clark), 311

Clusin, William T. (See Mason et al.), 633 COBB, LEONARD A. Prehospital cardiac care: Does it make a

difference, 316 Cody, Robert J., Franklin, Kenneth W., and Laragh, John H. Postural hypotension during tilt with chronic capto-pril and diuretic therapy of severe congestive heart failure, 480

COHEN, LAWRENCE S. (See Lewis et al.), 498
COHN, JAY N. Vasodilator therapy: Implications in acute myocardial infarction and congestive heart failure, 773

Collinson, Hector K. (See D'Cruz et al.), 295
Colsen, Peter R. (See Davis et al.), 289
Come, Patricia C., Isaacs, Randi E., and Riley, Marilyn F.
Diagnostic accuracy of M-mode echocardiography in
active infective endocarditis and prognostic implications of ultrasound-detectable vegetation, 839

CONETTA, DONALD A. (See Geiser, Skorton, and Conetta), 905—. (See Geiser et al.), 1056

Conti, C. Richard (See Whittle et al.), 161

—, Feldman, Robert L., and Pepine, Carl J. Coronary artery spasm: Prevalence, clinical significance, and provoca-

spasm: Prevalence, clinical significance, and provocative testing, 584

—. (See Geiser et al.), 1056
COOK, CHARLES A. (See Waeber et al.), 384
COOLEY, DENTON A. (See Gillette et al.), 66
CORBETT, JAMES C. (See Nicod et al.), 822
COVITZ, WESLEY, MOORE, H. VICTOR GRAY, BONNIE, BROWN, MARK, AND STRONG, WILLIAM B. Assessment of Fontan graft protection by the dispayable preference pulments again. patency by radionuclide perfusion pulmonary scan in tricuspid atresia with previous Glenn shunt, 1072 (Brief comm.)

CRAIGE, E. (See Sheps et al.), 519 CRUICKSHANK, JOHN M. (See Ramsdale et al.), 459

CURIN, T. (See Pfisterer et al.), 92 CURIONE, MARIO, LATINI, SANDRA, FERRANTI, EDOARDO, AND PULET-TI, MARIO. M-mode echogram in differential diagnosis of VT vs SVT with aberration, 1084 (Letter to Editor)

CURRY, R. CHARLES (See Whittle et al.), 161

Davidson, Robert. (See Schiff et al.), 562
Davies, Ross A. (See Lewis et al.), 498
Davis, J. C. (See Goldberg et al.), 75
Davis, Martin D., Caspi, Avi, Lewis, Basil S., Milner, Selwyn,
Colsen, Peter R., and Barlow, John B. Two-dimen-

b'Cruz, Ivan A., Collinson, Hector K., Gerrardo, Leo, and Hensel, Priscilla. Two-dimensional echocardiographic features of submitral left ventricular aneurysm, 289 (Brief comm.)

D'Cruz, Ivan A., Collinson, Hector K., Gerrardo, Leo, and Hensel, Priscilla. Two-dimensional echocardiographic detection of staphylococcal vegetation of the body of the collision of th attached to calcified mitral anulus, 295 comm.)

DEL NEGRO, ALBERT A. (See Waller et al.), 1071
DEMOS, TERRENCE C. (See Moncada et al.), 263
DESAI, RAJENDRA. (See Nair et al.), 226
DEVEREUX, RICHARD B. (See Reichek and Devereux), 202
—, AND SACHS, IRENE. Detection of abnormal LV function in

mitral regurgitation by echographic mitral-septal separation, 1086 (Letter to Editor)
DE ZWANN, CHRIS, BAR, FRITS W. H. M., AND WELLENS, HEIN J. J.

Characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction, 730 JONATHAN R., AND ESTES, N. MARK. ECG changes associated with iatrogenic left pneumothorax simulat-

DIAMOND, ing anterior myocardial infarction, 303 (Brief comm.)

DIETRICH, PETER. (See Lui et al.), 567

District, Peter. (See Lui et al.), 507

— (See Lee et al.), 1076

Diosi, Peter. (See Georgescu et al.), 572

Doi, Yoshinori, Sugiura, Tetsuro, Bishop, Richard L., Paladino, Delores, Moreau, Kathleen, and Spodick, David H. High-speed echophonocardiographic detection of tricuspid valve prolapse in mitral valve prolapse with discrepancy in onset of systolic murmur, 301 (Brief comm.)

Dorr, Rolf. (See Meyer et al.), 132 Dorsey, Thomas J. (See Singer, Rowen, and Dorsey), 131 Dressendorfer, Ruddleh H., Smith, Joan L., Amsterdam, Ezra A., and Mason, Dean T. Reduction of submaximal exercise myocardial oxygen demand post-walk training program in coronary patients due to improved physical work efficiency, 358

Dudell, Golde G. (See Valdes-Cruz et al.), 505
Duff, Henry J. (See Woosley et al.), 737
Duffy, C. Eliss (See Swiryn et al.), 168
Dwyer, Edward M., Jr., Pepe, Anthony J., and Pinkernell,
Bruce H. Effects of beta-adrenergic blockade with pindolol versus placebo in coronary patients with stable angina pectoris, 830 Dwyer, Richard M. (See Lee et al.), 1076

Effert, Sven. (See Meyer et al.), 132 Effron, Mark. (See Mehlman et al.), 137 Eisner, David R. (See Friedman et al.), 194

ELLIOTT, LARRY P. (See Rogers et al.), 185 ENGLE, W. KING. (See Gottdiener et al), 525 ENGLERT, MARC. (See Vandenbossche and Englert), 922 EPSTEIN, STEPHEN E. (See McManus et al.), 451 (See Sheehan and Epstein, 973 ESTES, N. MARK (See Diamond and Estes), 303 ESTIOKO, MANUEL. (See Kay, Estioko, and Wiener), 338

Falk, Rodney H. (See Zelinger, Falk, and Hood), 1073
FALKOFF, MICHAEL. (See Baran et al.), 1066
FARAGHER, ERIC B. (See Ramsdale et al.), 459
FELDMAN, ROBERT L. (See Whittle et al.), 161
—. (See Conti, Feldman, and Pepine), 584
FERRANTI, EDOARDO. (See Curione et al.), 1081

FIRTH, BRIAN G. (See Johnson et al.), 4
FISHER, JEFFREY, SONNENBLICK, EDMUND H., AND KIRK, EDWARD S.
Increased ventricular fibrillation threshold with severe myocardial ischemia, 966

FORFAR, J. COLIN. Prediction of hemorrhage during long-term oral coumarin anticoagulation by excessive prothrombin

ratio, 445 (Brief comm.)

FORKER, ALAN D. (See Marshall and Forker), 401

FORST, DAVID. (See O'Rourke, Badke, and Forst), 489

FOSTER, CARL. (See Pollock et al.), 363

Francis, Gary S., Sharma, Bim, and Hodges, Morrison. Comparative hemodynamic effects of dopamine and dobuta-

mine in patients with acute cardiogenic circulatory collapse, 995

FRANKLIN, KENNETH W. (See Cody, Franklin, and Laragh), 480

FRIEDMAN, MARK J., SAHN, DAVID J., GOLDMAN, STEVEN, EISNER, DAVID R., GITTINGER, NORMAN C., LEDERMAN, FRANK L., PUCKETTE, CHARLES M., AND TIEMANN, JEROME J. High predictive accuracy for detection of left main coronary artery disease by antilog signal processing of two-dimensional echocardiographic images, 194

FUKUYAMA, TAKAYA. (See Shimokawa et al.), 142 Furey, Warren. (See Mehlman et al.), 137

GALLAGHER, JOHN J. (See Baran et al.), 1066

GANG, E. (See Goldberg et al.), 75 GARRETT, BRUCE N., AND KAPLAN, NORMAN M. Efficacy of slow infusion of diazoxide in the treatment of severe hyper-

tension without organ hypoperfusion, 390 GARSON, ARTHUR, JR. (See Gillette et al.), 66 GARSON, ARTHUR, JR. (See Gillette et al.), 66

— (See Beder et al.), 848
GASSNER, SIDNEY. (See Korczyn et al.), 443
GAUM, WINSTON E. (See Biancaniello et al.), 852
GAVRAS, HARALAMBOS P. (See Waeber et al.), 384
GAVRAS, IRRNE. (See Waeber et al.), 384
GEISER, EDWARD A., SKORTON, DAVID J., AND CONETTA, DONALD A.
Quantification of left ventricular function by two-dimensional polescoparidownshus Consideration of the

dimensional echocardiography: Consideration of fac-

 —, Ariet, Mario, Conetta, Donald A., Lupkiewicz, Stefan M.,
Christie, Leonard G., Jr., and Conti C. Richard.
Dynamic three-dimensional echocardiographic reconstruction of the intact human left ventricle: Technique and initial observations in patients, 1056

Georgescu, Leonida, Diosi, Peter, Plavosin, Livia, Butiu, Ioan, and Herzog, Georgeta. Balkan nephritis and adenovirus infection, 572 (Letter to Editor) GERRARDO, LEO. (See D'Cruz et al.), 295

GETTES, LEONARD S., HILL, JEFFREY L., SAITO, TOMOAKI, AND KAGIYAMA, YUTAKA. Factors related to vulnerability to

arrhythmias in acute myocardial infarction, 667
GILBERT, BARRY K. (See Wood et al.), 649
GILLETTE, PAUL C., GARSON, ARTHUR, JR., COOLEY, DENTON A., AND
MCNAMARA, DAN G. Prolonged and decremental antegrade conduction properties in right anterior accessory connections: Wide QRS antidromic tachycardia of left bundle branch block pattern without Wolff-Parkinson-White configuration in sinus rhythm, 66 . (See Hesslein et al.), 351

 . (See Rossi et al.), 415
 . (See Beder et al.), 848 (See Karpawich et al.), 1001 GILLUM, RICHARD F., PRINEAS, RONALD J., AND KEBEDE, JOSEPH. Independent effects of Na reduction and K supplementation on blood pressure, 449 (Letter to Editor)

-. Will persons on low-sodium diets become iodine deficient,

1084 (Letter to Editor) GINSBURG, ROBERT, LAMB, IRENE H., SCHROEDER, JOHN S., HU,
MARIE, AND HARRISON, DONALD C. Randomized doubleblind comparison of nifedipine and isosorbide dinitrate therapy in variant angina pectoris due to coronary artery spasm, 44

AND BRISTOW, MICHAEL R. Reply, 928 (Letter to Editor) GIROD, DONALD A. (See Hurwitz et al.), 421 GIROTTI, ARNOLDO L. (See Rutitzky, Girotti, and Rosenbaum),

GITTINGER, NORMAN C. (See Friedman et al.), 194 GLAGOV, SEYMOUR. (See Mehlman et al.), 145

Glogar, Dietmar. (See Rude et al.), 986
Gold, Herman K. (See Leinbach and Gold), 768
Goldberg, D., Reiffel, J. A., Davis, J. C., Gang, E., Livelli, F.,
And Bigger, J. T., Jr. Electrophysiologic effects of
procainamide on sinus function in patients with and

without sinus node disease, 75
Goldberg, Leon I., and Raifer, Sol I. Sympathomimetic amines: Potential clinical applications in ischemic heart disase, 724

GOLDBERG, STANLEY J. (See Valdes-Cruz et al.), 505
—, KECECIOGLU-DRAELOS, ZOE, SAHN, DAVID J., VALDES-CRUZ, LIL-LIAM M., AND ALLEN, HUGH D. Range gated echo-Doppler velocity and turbulence mapping in patients with valvular aortic stenosis, 858
GOLDMAN, STEVEN (See Friedman et al.), 194

(See Morrison et al.), 1081

GOLDSTEIN, SIDNEY. The necessity of a uniform definition of sudden coronary death: Witnessed death within 1 hour of the onset of acute symptoms, 156

GOOCH, ALDEN S. Downward displacement of competent aortic valve prosthesis without left ventricular enlargement: A new sign of ascending root dissection in absence of valvular regurgitation, 920 (Brief comm.)

GORDON, TAVIA. Recent decline in coronary disease mortality in the United States: Part of a general decline in mortal-ity, 151 (Letter to Editor)

 AND KANNEL, WILLIAM B. Multiple risk functions for predicting coronary heart disease: The concept, accuracy, and application, 1031

GORLIN, RICHARD. Role of coronary vasospasm in the pathogeneis of myocardial ischemia and angina pectoris, 598 GOTTDIENER, JOHN S., HAWLEY, ROLLIN J., MARON, BARRY J., BERTORINI, TULIO F., AND ENGLE, W. KING. Character-

istics of the cardiac hypertrophy in Friedreich's ataxia, 525 Graboys, Thomas B. The stampede to stimulation-numerators

and denominators revisited relative to electrophysio-logic study of ventricular arrhythmias, 1089

Graham, Thomas P., Jr. (See Artman and Graham), 1040 Gray, Bonnie. (See Covitz et al.), 1072

GREEN, KENNETH G. The role of hypertension and downward changes of blood pressure in the genesis of coronary atherosclerosis and acute myocardial ischemic attacks,

GREENSPAN, ALLAN M. (See Horowitz et al.), 604 GRIFFIN, JERRY C. (See Mason et al.), 633

GRUENTZIG, ANDREAS. Results from coronary angioplasty and implications for the future, 779

GUERRERO, JUAN. (See Nademanee et al.), 802

— (See Josephson et al.), 978

GUMBINER, CARL H. (See Karpawich et al.), 1001

GUNNAR, ROLF M. (See O'Connell et al.), 85

— (See Moncada et al.), 263 — (See LaBarre, O'Connell, and Gunnar), 308 GJTGESELL, HOWARD P. (See Hesslein et al.), 351

Hackshaw, Barry T. (See Smuckler et al.), 897 Hager, W. David. (See Morrison et al.), 1081 Haiat, Robert. Second comment, 927 (Letter to Editor) HALE, DAVID. (See Moncada et al.), 263 HALL, ROBERT J. (See Rossi et al.), 415

- HAMER, ANGAS, PETER, THOMAS, AND MANDEL, WILLIAM J. Reply, 306 (Letter to Editor)
- Hannington-Kiff, J. G. Hyperadrenergic-effected limb causalgia: Relief by IV pharmacologic norepinephrine blockade,
- 152 (Letter to Editor)
 HANSEN, HERBERT. (See Schön et al.), 532

 —. (See Schön et al.), 548

- —, (See Schon et al.), 548

 HARKEN, ALDEN H. (See Martin et al.), 960

 HARRIS, LOWELL D. (See Wood et al.), 649

 HARRISON, DONALD C. (See Ginsburg et al.), 44

 HARTZ, RENEE. (See Mehrnan et al.), 137

 HAUTVAST, J. G. A. J. (See Knuiman, West, and Hautvast), 447

 HAWLEY, ROLLIN J. (See Gottdiener et al.), 525
- HAWLEY, ROLLIN J. (See Gottalener et al.), 526
 HAY, RICK. (See Mehlman et al.), 145
 HECHT, HARVEY S., AND ZIBELLI, LOUIS R. First-pass radionuclide angiography in visualization of acute myocardial infarction ventricular septal defect and assessment of ventricular function, 436 (Brief comm.)
- (See Josephson et al.), 978
 (See Josephson et al.), 1008
 Heger, James J. (See Rinkenberger et al.), 177
 (See Prystowsky et al.), 426
- (See Zipes, Prystowsky, and Heger), 610 (See Zipes et al.), 789

- (See Lipse et al.), 163

 HELIN, M. (See Raumio et al.), 32

 HELLMAN, CHARLES. (See Pollock et al.), 363

 HENDERSON, GENE. (See Nicod et al.), 822

 (See Nademanee et al.), 950

 (See Nademanee et al.), 950
- HENDRY, P. (See Biro, Beresford-Kroeger, and Hendry), 870

- HENDRY, F. (See Bird, Berestord-Kroeger, and Hendry), 870
 HENSEL, PRISCILLA. (See D'Cruz et al.), 295
 HENZE, EBERHARD. (See Schelbert et al.), 588
 HERMANOVICH, JOHN. (See Awan et al.), 474
 HERZOG, GEORGETA. (See Georgescu et al.), 572
 HESSLEIN, PETER S., GUTGESELL, HOWARD P., GILLETTE, PAUL C.,
 AND MCNAMARA, DAN G. Exercise assessment of sinoatrial node function following the Mustard operation,
- HEUSER, RICHARD R., ACHUFF, STEPHEN C., AND BRINKER, JEFFREY
 A. Inadvertent division of an anomalous left anterior descending coronary artery during complete repair of tetralogy of Fallot: 22-year follow-up, 430 (Brief comm.)
- Hill, Jeffrey L. (See Gettes et al.), 667 Hills, L. David. (See Johnson et al.), 4 Hirata, Tsuneo. (See Imaizumi et al.), 887
- HIRAH, TSUNEO. (See Imazumi et al., 867 HIRSH, STANTON A. Acute allergic reaction with coronary vaso-spasm, 928 (Letter to Editor) HODGES, MORRISON. (See Francis, Sharma, and Hodges), 995 HOOD, WILLIAM B., JR. (See Zellinger, Falk, and Hood), 1073 HOPKINS, JAMES. (See Josephson et al.), 978
- (See Josephson et al.), 1008
- HOROWITZ, LEONARD N., SPIELMAN, SCOTT R., GREENSPAN, ALLAN M. Role of programmed stimulation in assessing vulnerability to ventricular arrhythmias, 604 (See Martin et al.), 960
- HOROWITZ, RICHARD S., AND MORGANROTH, JOEL. Immediate detection of early high-risk patients with acute myocardial infarction using two-dimensional echocardiographic evaluation of left ventricular regional wall motion abnormalities, 814
- Hu, Marie. (See Ginsburg et al.), 44 Huang, Henry. (See Schön et al.), 548
- HUANG, SUNG-CHENG. (See Schon et al.), 548
 HUANG, SUNG-CHENG. (See Schon et al.), 532
 HURWITZ, ROGER A., TREVES, SALVADOR, KEANE, JOHN F., GIROD,
 DONALD A., CALDWELL, RANDALL L. Current value of
 radionuclide angiocardiography for shunt quantification and management in patients with secundum atrial septal defect, 421
- HUSSEIN, HANY. (See Lee et al.), 1076 HUYSMANS, FRANS TH.M., THIEN, THEO A., AND KOENE, ROBERT A. Combined intravenous administration of diazoxide and beta-blocking agent in acute treatment of severe hypertension or hypertensive crisis, 395
- Ikeda, Richard M. (See Lee et al.), 1076 Imaizumi, Tsutomu, Orita, Yasuhiko, Koiwaya, Yasushi, Hirata, Tsuneo, and Nakamura, Motoomi. Utility of two-dimensional echocardiography in the differential diagnosis of the etiology of aortic regurgitation, 887

- INAGAKI, YOSHIAKI. (See Masuda et al.), 230 INTARACHOT, VANIDA. (See Nademanee et al.), 802 ISAACS, RANDI E. (See Come, Isaacs, and Riley), 839
- Jackman, Warren M. (See Rinkenberger et al.), 177 —. (See Prystowsky et al.), 426 —. (See Zipes et al.), 789
- JAMES, D. GERAINT, AND THOMSON, A. Recognition of the diverse cardiovascular manifestations in Behcet's disease,

- JANATUINEN, E. (See Raunio et al.), 32
 JANICKI, JOSEPH S. (See Weber, Andrews, and Janicki), 639
 JOHNSON, SARAH A. (See O'Connell et al.), 85
 JOHNSON, STACEY M., MAURITSON, DAVID R., WINNIFORD, MICHAEL
 D., WILLERSON, JAMES T., FIRTH, BRIAN G., CAREY, JOAN
 R., AND HILLIS, L. DAVID. Continuous electrocardiographic monitoring in patients with unstable angina
 pectoris: Identification of high-risk subgroup with
 severe coronary disease, variant angina, and/or
 impaired early prognesis.
- Jones, Michael. (See McManus et al.), 451
 Josephson, Mark E. (See Horowitz et al.), 604

 —. (See Martin et al.), 960
 Josephson, Martin A., Hecht, Harvey S., Hopkins, James, Guer-
 - RERO, JUAN, AND SINGH, BRAMAH N. Comparative effects of oral verapamil and propranolol on exercise-induced myocardial ischemia and energetics in patients with coronary artery disease: Single-blind placebo cross-over evaluation using radionuclide ventriculography,
- -, Brown, B. Greg., Hecht, Harvey S., Hopkins, James, Pierce, Cynthia D., and Petersen, Robert B. Noninvasive detection and localization of coronary stenoses in patients: Comparison of resting dipyridamole and exercise thallium-201 myocardial perfusion imaging,
- JUTZY, KENNETH R. (See Mitchell et al.), 57

- KAGIYAMA, YUTAKA. (See Gettes et al.), 667
 KAIMAL, KRISHNA P. Computed tomography in the diagnosis of pericardial cyst, 566 (Brief comm.)
 KALF, VICTOR. (See Rizi et al.), 292
 KANAIDE, HIDEO. (See Kaseda et al.), 569
 KANNAN, RAMASWAMY. (See Nademanee et al.), 950
 KANNAL, WILLIAM B. (See Gordon and Kannel), 1031
 KAPLAN, NORMAN M. (See Garrett and Kaplan), 390
 KAPLAN, SAMUEL. (See Biancaniello et al.), 852
 KARAFFA ASTEPHANIE. (See Rude et al.), 986

- KARAFFA, STEPHANIE. (See Rude et al.), 986
- KARPAWICH, PETER P., GUMBINER, CARL H., GILLETTE, PAUL C., Shih, Joung-Yi, Zinner, Alex, and Lewis, Robert. Comparative electrophysiologic effects of digoxin in the non-sedated chronically instrumented puppy, 1001
- Kaseda, Shunichi, Koiwaya, Yasushi, Tajimi, Tsukasa, Mitsutake, Arahito, Kanaide, Hideo, Takeshita, Akira, Kikuchi, Yutaka, Nakamura, Motoomi, Mayumi, Hisanohi, Komori, Masataka, and Tokunaga, Kouichi. Huge false aneurysm due to rupture of the right coronary artery in Behcet's syndrome, 569 (Brief comm.)
- KATO, YASUMICHI. (See Kinoshita et al.), 1090 KAVEY, RAE-ELLEN WEBB, BLACKMAN, MARIE S., AND SONDHEIMER, HENRY M. Incidence and severity of chronic ventricu-lar dysrhythmias after repair of tetralogy of Fallot,
- KAWASAKI, TAKESHI. (See Kinoshita et al.), 1090 KAY, RICHARD, BLAKE, JEFFREY AND RUBIN, DAVID. Possible coronary spasm rebound to abrupt nifedipine withdrawal, 308 (Letter to Editor)

 —, ESTIOKO, MANUEL, AND WIENER, ISAAC. Primary sick sinus syndrome as an indication for chronic pacemaker
- therapy in young adults: Incidence, clinical features, and long-term evaluation, 338
- Keane, John F. (See Hurwitz et al.), 421 Kebede, Joseph. (See Gillum, Prineas, and Kebede), 449 Kececioglu-Draelos, Zoe. (See Goldberg et al.), 858

KEEFE, DEBORAH L., PETERS, FLORA, AND WINKLE, ROGER A. Randomized double-blind placebo controlled crossover trial documenting oral lorcainide efficacy in suppression of symptomatic ventricular tachyarrhythmias, 511

—. (See Mason et al.), 633 KEELAN, MICHAEL H., Jr. (See Olinger, Thompson, and Keelan), 440

KELLER, MARTIN D. (See Lewis et al.), 123
KESSLER, KENDISTH M., KIEVAL, JOSHUA, SAKSENA, SANJEEV,
SANDERSON, TAMARA L., AND MYERBURG, ROBERT J.
Echographic features of posterior left ventricular wall
pseudoaneurysm due to Escherichia coli endocarditis, 139 Brief comm.)

—. (See Myerburg et al.), 615
KHAN, ABDUL H. Postmyocardial infarction Dressler's syndrome,
926 (Letter to Editor)
KHURI, SHUKRI F. (See Rude et al.), 986
KIEVAL, JOSHUA. (See Kessler et al.), 139

KIKUCHI, YUTAKA. (See Kaseda et al.), 569

KINOSHITA, SHINJI, KATO, YASUMICHI, KAWASAKI, TAKESHI, AND OKIMORI, KATSUSHI. Ventricular tachycardia initiated by late-coupled ventricular extrasystoles: The concept of longitudinal dissociation in the microreentry pathway, 1090

KINSEY, JAMES H. (See Wood et al.), 649
KIRK, EDWARD S., AND SONNENBLICK, EDMUND H. Newer concepts in the pathophysiology of ischemic heart disease,

. (See Fisher, Sonnenblick, and Kirk), 966 KISSLO, JOSEH A. (See Bardy et al.), 1068 KITAZUME, HIDEMASA. (See Kramer et al.), 283 KIZAKEVICH, P. N. (See Sheps et al.), 519 KLINE, ROBERT C. (See Rizi et al.), 292

KLINER, ROBERT A. (See Rude et al.), 986 KNUIMAN, J. T., WEST, C. E., AND HAUTVAST, J. G. A. J. Role of diet to HDL-cholesterol and coronary disease: Serum total and HDL-cholesterol in boys and men from develop-ing and developed countries, 447 (Letter to Editor)

Kobrin, I. (See Yagil et al.), 310 Koene, Robert A. P. (See Huysmans, Thien, and Koene), 395

Koiwaya, Yasushi. (See Shimokawa et al.), 142 —. (See Kaseda et al.), 569

- (See Raseda et al.), 509
- (See Inaizumi et al.), 887
- (See Matsuguchi et al.), 918
KOMORI, MASATAKA. (See Kaseda et al.), 569
KORCZYN, AMOS D., SHAVIT, SARA, GASSNER, SIDNEY, AND VIDNE,
BERNARDO. Denervation supersensitivity in the canine heart: Catecholamine-enhanced positive chronotropy without altered arrhythmogenicity, 443 (Brief comm.)

Kossowsky, Warren A., Lyon, Alan F., and Spain, David M. Reply, 926 (Letter to Editor)
Kramer, John R. (See Mellino et al.), 222

—, Kitazume, Hidemasa, Produpitt, William L., and Sones, F. Mason, Jr. Clinical significance of isolated coronary bridges: Benign and frequent condition involving the left anterior descending artery, 283
Kuhl, David E. (See Schön et al.), 532

(See Schölbert et al.), 538

(See Schelbert et al.), 588

Labadie, Enrique. (See Morrison et al.), 1081

LaBarre, Terry R., O'Connell, John B., and Gunnar, Rolf M.

Captopril therapy for severe CHF: Hypotensive response in presence of markedly elevated PRA, 308 (Letter to Editor)

L'ABBATE, A. (See Biagini et al.), 13
LAM, WILFRED. (See Swiryn et al.), 319
—. (See Strasberg et al.), 795
LAMB, IRENE H. (See Ginsburg et al.), 44

LAMBERTZ, Heinz. (See Meyer et al.), 132 LANESE, RICHARD R. (See Lewis et al.), 123 LANGFORD, HERBERT G. Reduced stroke incidence in HDFP enhanced by strong support system, 1087 (Letter to

LARAGH, JOHN H. (See Cody, Franklin, and Laregh), 480 LATINI, SANDRO. (See Curione et al.), 1081 LEDERMAN, FRANK L. (See Friedman et al.), 194

LEE, GARRETT. (See Lui et al.), 567

 KEDA, RICHARD M., DWYER, RICHARD M., HUSSEIN, HANY, DIETRICH, PETER, AND MASON, DEAN T. Feasibility of intravascular laser irradiation for in vivo visualization and therapy of cardiocirculatory diseases, 1076 (Brief

comm.)
Leibel, B. (See Yagil et al.), 310
Leinbach, Robert C., and Gold, Herman K. Coronary angiography during acute myocardial infarction: A search for spasm, 768

LEVENSON, J. A. (See Payen et al.), 379 LEVER, HARRY M. (See Mellino et al.), 222 LEVINE, HEREFET J. Introduction, 583 LEWIS, BASIL S. (See Davis et al.), 289

Lewis, Fighard P., Lanese, Richard R., Stang, John M., Chiri-kos, Tromas N., Keller, Martin D., and Warren, James V. Reduction of mortality from prehospital myocardial infarction by prudent patient activation of

mobile coronary care system, 123
Lewis, Robert. (See Karpawich et al.), 1001
Lewis, Samuel E. (See Nicod et al.), 822
Lewis, Sandra J. (See Mitchell et al.), 57

Lewis, Sandra J. (See Mitchell et al.), 57

Lewis, Stanley M., Riba, Arthur L., Berger, Harvey J., Davies, Ross A., Wackers, Frans J. Th., Alexander, Jona-than, Sands, Milton J., Cohen, Lawrence S., and Zaret, Barry L. Radionuclide angiographic exercise left ventricular performance in chronic aortic regurgitation: Relationship to resting echographic ventricular dimensions and systolic wall stress index, 498 Lie, J. T. Aortic dissection in Turner's syndrome, 1077 (Brief

comm.)

LINDEMANN, JON P., BAILEY, JOHN C., AND WATANABE, AUGUST M. Potential biochemical mechanisms for regulation of the slow inward current: Theoretical basis for drug action, 746 Linnerup, A. C. (See Pollock et al.), 363 Lipton, Martin J. (See Brundage and Lipton), 313

LIVELLI, F. (See Goldberg et al.), 75 LOMBARDI, FEDERICO. (See Malliani and Lombardi), 575 LORELL, BEVERLY. (See Smith et al.), 716

LOVE, LEON. (See Moncada et al.), 263 Low, REGINALD I. (See Lui et al.), 567 Lown, Bernard. Management of patients at high risk of sudden

death, 689 LUETZELER, JOSEF. (See McManus, Luetzeler, and Roberts), 298 LUI, HENRY K., LEE, GARRETT, DIETRICH, PETER, LOW, REGINALD I., AND MASON, DEAN T. Flecainide-induced QT prolongation and ventricular tachycardia, 567 (Brief comm.)

LULL, ROBERT J. (See Sobol et al.), 563 LUPKIEWICZ, STEFAN M. (See Geiser et al.), 1056 LYON, ALAN F. (See Kossowsky, Lyon, and Spain), 926 LYONS, KENNETH P. (See Butman et al.), 912

M

MALLIANI, ALBERTO, AND LOMBARDI, FEDERICO. Consideration of the fundamental mechanisms eliciting cardiac pain, 575 Mandel, William J. (See Hamer, Peter, and Mandel), 306

MANTLE, JOHN A. (See Rackley et al.), 154

—. Regular insulin dose in GIK formulation for acute myocardial

MARON, BARRY J. (See Gottdiener et al.), 525
MARSH, JAMES D. (See Smith et al.), 716

MARSH, JAMES D. (See Smith et al.), 716

MARSHALL, JOHN B., AND FORKER, ALAN D. Cardiovascular effects of tricyclic antidepressant drugs: Therapeutic usage, overdose, and management of complications, 401

MARTIN, JACK L., UNTEREKER, WILLIAM J., HARKEN, ALDEN H., HOROWITZ, LEONARD N., AND JOSEPHSON, MARK E. Aneurysmectomy and endocardial resection for ventricular tachycardia: Favorable hemodynamic and antionhythmic accounts in the debal left. antiarrhythmic results in patients with global left ventricular dysfunction, 960

MASERI, A. (See Biagini et al.), 13

Mason, Dean T. Progress report on publication goals, 1

—. (See Vera, Awan, and Mason), 80

—. (See Dressendorfer et al.), 358

(See Awan et al.), 474 (See Lui et al.), 567

(See Riemenschneider et al.), 834

(See Wikman-Coffelt, Parmley, and Mason), 934

(See Lee et al.), 1076

Mason, Jay W. (See Mitchell et al.), 57

—, Swerdlow, Charles D., Winkle, Roger A., Griffin, Jerry C.,
Ross, David L., Keefe, Deborah L., and Clusin,
William T. Programmed ventricular stimulation in predicting vulnerability to ventricular arrhythmias and their response to antiarrhythmic therapy, 633 Massumi, All. (See Rossi et al.), 415

MASSUM, ALL (See Rossi et al.), 413

MASUDA, YOSHIAKI, YOSHIDA, HIDBO, MOROOKA, NOBUHIRO, TAKAHASHI, OSHIAKI, WATANABE, SHIGERU, INAGAKI, YOSHIAKI, UCHIYAMA, GYO AND TATENO, YOKIO. ECG synchronized computer tomography in clinical evaluation of total and regional cardiac motion: Comparison of postmyocardial infarction to normal hearts by rapid sequential imaging, 230

Matsuguchi, Takeyuki. (See Shimokawa et al.), 142

-, Koiwaya, Yasushi, Nakagaki, Osamu, Orita, Yasuhiko, and Nakamura, Motoomi. Negative U wages and peaked T waves without ST changes during spontaneous and ergonovine-induced vasospastic angina, 918 (Brief

MATTSON, SÖREN. (See Ulmsten, Bernstein, and Mattson), 448
MAURER, GERALD, AND NANDA, NAVIN C. Two-dimensional echocardiographic identification of intracardiac leiomyomatosis, 915 (Brief comm.)

MAURITSON, DAVID R. (See Johnson et al.), 4 MAYUMI, HISANORI. (See Kaseda et al.), 569

MAZZEI, M. G. (See Biagini et al.), 13 McGrath, Francis J. (See Waller et al.), 1071 McManus, Bruce M., Luetzeler, Josef and Roberts, William C. Total anomalous pulmonary venous connection: Survival for 62 years without surgical intervention, 298 (Brief comm.)

—, Waller, Bruce F., Jones, Michael, Epstein, Stephen E., and Roberts, William C. The case for preoperative coro-nary angiography in patients with tetralogy of Fallot and other complex congenital heart diseases, 451

McNamara, Dan G. (See Gillette et al.), 66 (See Beder, Nihill, and McNamara), 134

(See Hesslein et al.), 351

— (See Beder et al.), 348

McWilliams, Gregory J. (See Besley et al.), 332

Mehlman, David J., Furry, Warren, Phair, John, Effron, Mark, and Hartz, Rener. Two-dimensional echocardiographic features diagnostic of isolated pulmonic valve endocarditis, 137 (Brief comm.)

—, Troncoso, Patricia, Hay, Rick, And Gladov, Seymour. Midsystolic click accompanying isolated bicuspid pulmonic valve, 145 (Brief comm.)

MEHTA, JAWAHAR, AND MEHTA, PAULETTE. Comparison of platelet function during exercise in normal subjects and coronary artery disease patients: Potential role of platelet activation in myocardial ischemia, 49

MEHTA, PAULETTE. (See Mehta and Mehta), 49

MELLINO, MARCELLO, SALCEDO, ERNESTO E., LEVER, HARRY M.,

VASUDEVAN, GOPALAN, AND KRAMER, JOHN R. Echo-graphic-quantified severity of mitral anulus calcification: Prognostic correlation to related hemodynamic, valvular, rhythm, and conduction abnormalities, 222

Merx, Wolfgang, (See Meyer et al.), 132
Meyer, Jürgen, Merx, Wolfgang, Dörr, Rolf, Lambertz, Heinz,
Bethge, Christian and Effert, Sven. Successful
treatment of acute myocardial infarction shock by

combined percutaneous transluminal coronary recan-alization (PTCR) and percutaneous transluminal cor-onary angioplasty (PTCA), 132 (Brief comm.)

MEYER, RICHARD A. (See Biancaniello et al.), 852
MICHELASSI, C. (See Biagini et al.), 13
MILNER, SELWYN. (See Davis et al.), 289
MIROWSKI, M., REID, PHILIP R., MOWER, MORTON M., AND WATKINS,
LEVI. Successful conversion of out-of-hospital lifethreatening arrhythmias with the implanted automat-

ic defibrillator, 147 (Brief comm.)

MITCHELL, L. Brent, Jutzy, Kennerth R., Lewis, Sandra J.,
Schroeder, John S., and Mason, Jay W. Intracardiac
electrophysiologic study of intravenous diltiazem and

MITSUTAKE, ARAHITO. (See Kaseda et al.), 569
MOFFATT, ROY. (See Schiff et al.), 438
—. (See Schiff et al.), 438
MOHUDDIN, SYED M. (See Nair et al.), 226

Moncada, Rogelio, Baker, Mark, Salinas, Matias, Demos, Ter-rence C., Churchill, Robert, Love, Leon, Reynes, Carlos, Hale, David, Cardoso, Manuel, Pifarre, ROQUE, AND GUNNAR, ROLF M. Diagnostic role of computed tomography in pericardial heart disease: Congenital defects, thickening, neoplasms, and effusions, 263

Moodie, Doulgas S. (See Besley et al.), 332 Moore, H. Victor. (See Covitz et al.), 1072

Moore, H. Victor. (See Covitz et al.), 1072
Moreau, Kathleen. (See Doi et al.), 301
Morganroth, Joel. (See Horowitz and Morganroth), 814
Morooka, Nobuhiro. (See Masuda et al.), 230
Morrison, Douglass, Goldman, Steven, Labadie, Enrique, and
Hager, W. David. Primary VF during EEG in a
psychiatric patient, 1081 (Letter to Editor)
Mower, Morton M. (See Mirowski et al.), 147
Muller, James E. (See Rude et al.), 986
Muller, Brann, J. (See Pfisterer et al.), 92
Muz, Jaroslaw. (See Wizenberg et al.), 468
Myerburg, Robert J. (See Kessler et al.), 139
—, Zaman, Liaquat, Kessler, Kenneth M., and Castellanos,

, Zaman, Liaquat, Kessler, Kenneth M., and Castellanos, Agustin. Evolving concepts of management of stable and potentially lethal arrhythmias, 615

NACCARELLI, GERALD V. (See Rinkenberger et al.), 177

— (See Prystowsky et al.), 426 — (See Zipes et al.), 789 Nademanee, Koonlawee, Singh, Bramah N., Guerrero, Juan, Hendrickson, JoAnn, Intarachot, Vanida, and Baky, SHERRIN. Accurate rapid compact analog method for the quantification of frequency and duration of myo-

cardial ischemia by semiautomated analysis of 24-hour Holter ECG recordings, 802

—, HENDRICKSON, JOANN, KANNAN, RAMASWAMY, AND SINGH, BRAMASWAMY, AND SINGH, BRAMAN N. Antiarrhythmic efficacy and electrophysiolog ic actions of amiodarone in patients with life-threatening ventricular arrhythmias: Potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of induced ventricular tachycardia, 950

NAIR, CHANDRA K., SKETCH, MICHAEL H., DESAI, RAJENDRA, MOHIUDDIN, SYED M., AND RUNCO, VINCENT. High prevalence of symptomatic bradyarrhythmias due to atrioventricular node-fascicular and sinus node-atrial dis-ease in patients with mitral anular calcification, 226 Najari, Ahmad. (See Schön et al.), 532

(See Schön et al.), 548 — (See Schön et al.), 548

NAKAGAKI, OSAMU. (See Matsuguchi et al.), 918

NAKAMOTO, SATORU. (See Chaignon et al.), 374

NAKAMURA, MOTOOMI. (See Shimokawa et al.), 142

— (See Kaseda et al.), 569

— (See Imaizumi et al.), 887

— (See Matsuguchi et al.), 918

NANDA, NAVIN C. (See Rosenzweig and Nanda), 435

— (See Maurer and Nanda), 915

... (See Maurer and Nanda), 915
... (See Ong, Nanda, and Barold), 917
... (See Baran et al.), 1066
Nanii, Amin A., and Blank, David. Increased 24-hour urine

Nanji, Amin A., and Blank, David. Increased 24-hour urine vanillylmandelic acid in patients without pheochromocytoma, 1088 (Letter to Editor)

Needham, Kathleen E. (See Awan et al.), 474

Newburger, Jane W. (See Borow and Newburger), 879

Newland, Hillary R. (See Ouzts and Newland), 928

Nicod, Pascal, Lewis, Samuel E., Corbett, James C., Buja, L. Maximilian, Henderson, Gener, Raskin, Philip, Rude, Robert E., and Willerson, James T. Increased incidence and clinical correlation of persistently abnormal technetium pyrophosphate myocardial scintirgrams technetium pyrophosphate myocardial scintigrams following acute myocardial infarction in patients with diabetes mellitus, 822

NIHILL, MICHAEL R. (See Beder, Nihill, and McNamara), 134 NIHOVANNOPOULOS, P. (See Venetis et al.), 924 NOMEIR, ABDEL-MOHSEN. (See Smuckler et al.), 897 NORRIS, ROBIN M. (See Abraham and Norris), 929

OATES, JOHN A. (See Woosley et al.), 737

O'CONNELL, JOHN B., WALLIS, DIANE, JOHNSON, SARAH A., PIFARRE, ROQUE, AND GUNNAR, ROLF M. Transient bundle branch block following use of hypothermic cardioplegia in coronary artery bypass surgery: High incidence without perioperative myocardial infarction, 85

(See LaBarre, O'Connell, and Gunnar), 308 OKADA, ROBERT D. (See Brown et al.), 1019

OKMORI, KATSUSHI. (See Kinoshita et al.), 1090
OKIMORI, KATSUSHI. (See Kinoshita et al.), 1090
OLINGER, GORDON N., THOMPSON, MELISH A., AND KEELAN, MICHAEL
H., JR. Optimal management of suspected thrombosis
of standard Björk-Shiley unmarked tilting disc mitral

valve prosthesis, 440 (Brief comm.) Olson, Harold G. (See Butman et al.), 912

ONG, LING S., NANDA, NAVIN C., AND BAROLD, S. SERGE. Twodimensional echocardiographic detection and diagnostic features of left ventricular papillary fibroelastoma, 917 (Brief comm.)

ORITA, YASUHIKO. (See Shimokawa et al.), 142

Orita, 1480Hiko. (See Shimokawa et al.), 142

—. (See Imaizumi et al.), 837

—. (See Matsuguchi et al.), 918
Ormiston, John A. (See Tei, Shah, and Ormiston), 1025
O'ROURKE, ROBERT A., BADKE, FREDERICK R., AND FORST, DAVID.
Comparative hemodynamic effects of digoxin vs nitroprusside in conscious dogs with aortocaval fistulainduced chronic left ventricular volume overload and

normal systolic performance, 489
OUZTS, HARVEY G., AND NEWLAND, HILLARY R. Cardiac arrest in pheochromocytoma, 928 (Letter to Editor)

Paladino, Delores. (See Doi et al.), 301 Palileo, Edwin A. (See Swiryn et al.), 168 —. (See Swiryn et al.), 319

(See Strasberg et al.), 795

-. (See Strasberg et al.), 150
-. (See Bauernfeind et al.), 941
PAPAPIETRO, SILVIO E. (See Rackley et al.), 154
PARMLEY, WILLIAM W., ROULEAU, JEAN-LUCIEN, AND CHATTERJEE,
KANU. Vasodilators in heart failure secondary to coronary artery disease, 625

— . (See Wikman-Coffelt, Parmley, and Mason), 934

PATEL, JIJIBHOY. (See Sobol et al.), 563

PAVEL, DAN. (See Swiryn et al.), 319

PAYEN, D. M., SAFAR, M. E., LEVENSON, J. A., TOTOMOKOUO, J. A.,
AND WEISS, Y. A. Prospective study of predictive factors determining borderline hypertensive individuals. als who develop sustained hypertension: Prognostic value of increased diastolic orthostatic blood pressure tilt-test response and subsequent weight gain, 379
PEARL, WILLIAM R. Massive lidocaine overdose, 1083 (Letter to

Editor)

PEPE, ANTHONY J. (See Dwyer, Pepe, and Pinkernell), 830

PEPINE, CARL J. (See Whittle et al.), 161

— (See Conti, Feldman, and Pepine), 584

PERLOFF, JOSEPH K. (See Child and Perloff), 1087

PETER, THOMAS. (See Hamer, Peter, and Mandel), 306 PETERS, FLORA. (See Keefe, Peters, and Winkle), 511

Peters, Flora. (See Reere, Peters, and Winkle), 511
Petersen, Robert B. (See Josephson et al.), 1008
Petrovick, M. L. (See Sheps et al.), 519
Petersen, M., Müller-Brand, J., Bründler, H., and Cueni, T.
Prevalence and significance of reversible radionuclide ischemic perfusion defects in symptomatic aortic valve disease patients with or without concomitant

PHAIR, JOHN. (See Mehlman et al.), 137
PHELPS, MICHAEL E. (See Schön et al.), 532

—. (See Schön et al.), 548

—. (See Schelbert et al.), 588
PIERCE, CYNTHIA D. (See Josephson et al.), 1008

PIFARRE, ROQUE. (See O'Connell et al.), 85

—. (See Moncada et al.), 263

PINKERNELL, BRUCE H. (See Dwyer, Pepe, and Pinkernell), 830

PIOMELLI, SERGIO. (See Valdes-Cruz et al.), 505

PITT, BERTRAM. (See Rizi et al.), 292

(See Popat and Pitt), 485
 AND THRALL, JAMES H. Evaluation of patients with ischemic heart disease by thallium-201 myocardial imaging,

PLAUTH, W. H., JR. (See Wilcox et al.), 148 PLAVOSIN, LIVIA. (See Georgescu et al.), 572 Pohost, Gerald. (See Brown et al.), 1019

POLLOCK, MICHAEL L., FOSTER, CARL, SCHMIDT, DONALD, HELLMAN, CHARLES, LINNERUD, A. C., AND WARD, ANN. Comparative analysis of physiologic responses to three different maximal graded exercise test protocols in healthy women, 363

POPAT, KISHOR D., AND PITT, BERTRAM. Hemodynamic effects of prostaglandin E, infusion in patients with acute myocardial infarction and left ventricular failure, 485

PRINEAS, RONALD J. (See Gillum, Prineas, and Kebede), 449

Princas, Ronald J. (See Kramer et al.), 283
Prystowsky, Eric N. (See Kramer et al.), 177

—, Heger, James J., Jackman, Warren M., Naccarelli, Gerald V., and Zipes, Douglas P. Post-myocardial infarction incessant supraventricular tachycardia due to con-cealed accessory pathway, 426 (Brief comm.)

— (See Zipes, Prystowsky, and Heger), 610

- (See Zipes, Prystowsky, and Fiegel, 510

- (See Zipes et al.), 789

Pracin, Michael J., Tresch, Donald D., Soin, Jagmeet S., and Brooks, Harold L. Evaluation of postresuscitation left ventricular global and segmental function by radionuclide ventriculography in sudden coronary death survivors of prehospital cardiac arrest: Correlation to subsequent short-term prognosis, 54
PUCKETTE, CHARLES M. (See Friedman et al.), 194

PULETTI, MARIO. (See Curione et al.), 1081

RABINOVITCH, MARK. (See Rizi et al.), 292

RACKLEY, CHARLES E., RUSSELL, RICHARD O., JR., ROGERS, WILLIAM J., MANTLE, JOHN A., AND PAPAPIETRO, SILVIO E. ULSTa-ble angina pectoris: Is it time to change our approach,

(See Rogers et al.), 185

RAHLLY, G. THOMAS, Jr. (See Zipes et al.), 789
RAIFER, SOL I. (See Goldberg and Raifer), 724
RAMSDALE, DAVID R., FARAGHER, ERIC B., BENNETT, DAVID H.,
BRAY, COLIN L., WARD, CHRISTOPHER, CRUICKSHANK,
JOHN M., YUSUF, SALIM pain relief in patients with acute myocardial infarc-

pain relief in patients with acute myocardial intarction by intravenous atenolol, 459

—, and Bennett, D. H. ASH in conditions other than hypertrophic cardiomyopathy, 1083 (Letter to Editor)

RASKIN, PHILIP. (See Nicod et al.), 822

RAUNIO, H., RISSANEN, V., HELIN, M., REHNBERG, S., ROMPPANEN, T., and JANATUINEN, E. Early pronounced ST segment depression with marked J point decline heralding acute lethal clinical myocardial infarction: Necropsyelectrographic correletive study, 32

electrocardiographic correlative study, 32 AND CLARK, T. J. H. Paroxysmal nocturnal dyspnea in CHF: Mechanisms and amniophylline therapy, 311

(Letter to Editor)
REHNBERG, S. (See Raunio et al.), 32
REICHEK, NATHANIEL, AND DEVEREUX, RICHARD B. Reliable estimation of peak left ventricular systolic pressure by M-mode echographic-determined end-diastolic relative wall thickness: Identification of severe valvular aortic stenosis in adult patients, 202

REID, PHILIP R. (See Mirowski et al.), 147 REIFFEL, J. A. (See Goldberg et al.), 75 REINECKE, CYNTHIA. (See Valdes-Cruz et al.), 505

REINES, CARLOS. (See Moncada et al.), 263 RIBA, ARTHUR L. (See Lewis et al.), 498 RIEMENSCHNEIDER, THOMAS A., BRENNER, ROBERT A., WIKMAN-COFFELT, JOAN, MASON, DEAN T. Maturational changes in cardiac muscle myosin adenosine triphosphatase activity relative to hemodynamic alterations in new-

born lambs, 834

Riley, Marilyn F. (See Come, Isaacs, and Riley), 839

Rinkenberger, Robert L., Prystowsky, Eric N., Jackman, Warren M., Naccarelli, Gerald V., Heger, James J., and REN M., NACCARELIJ, UERALD V., HEGER, JAMES J., AND ZIPES, DOUGLAS P. Drug conversion of nonsustained ventricular tachycardia to sustained ventricular tachycardia during serial electrophysiologic studies: Identification of drugs that exacerbate tachycardia and potential mechanism, 177

RISSANEN, V. (See Raunio et al.), 32

RITMAN, ERIK L. (See Wood et al.), 649

RIZI, H. ROGER, KLINE, ROBERT C., BESOZZI, MYRWOOD C., KALF,
VICTOR, RABINOVITCH, MARK, CHAN, WILLIAM, THRALL,
JAMES H., AND PITT, BERTRAM. Walk-through angina
phenomenon demonstrated by graded exercise radionuclide ventriculography: Possible coronary spasm
mechanisms, 292 (Brief comm.)
ROBB, RICHARD A. (See Wood et al.), 649
ROBERTS, WILLIAM C. Complications of cardiac valve replacement:
Characteristic abnormalities or prostheses pertaining
to any or specific site. 113

to any or specific site, 113

(See Siegel and Roberts), 210 (See McManus, Luetzeler, and Roberts), 298

(See McManus et al.), 451 No cardiac catheterization before cardiac valve replacement a mistake, 930

(See Waller et al.), 1071

ROBINSON, GERALD. (See Schön et al.), 532
RODEN, DAN M. (See Woosley et al.), 737
ROGERS, WILLIAM J. (See Rackley et al.), 154

—, SMITH, RICHARD L., BREAM, PETER R., ELLIOT, LARRY P.,
RACKLEY, CHARLES E., AND RUSSELL, RICHARD O., JR. Quantitative axial oblique contrast left ventriculography: Validation of the method by demonstrating improved visualization of regional wall motion and mitral valve function with accurate volume determi-

nations, 185 ROLNITZKY, LINDA M. (See Bigger, Weld, and Rolnitzky), 660

ROLINTZKY, LINDA M. (See Bigger, Weld, and ROMPPANEN, T. (See Raunio et al.), 32 ROSEN, KENNETH M. (See Swiryn et al.), 168 —. (See Swiryn et al.), 319 —. (See Strasberg et al.), 795 —. (See Bauernfeind et al.), 941

ROSENBAUM, MAURICIO B. (See Rutitzky, Girotti, and Rosenbaum), 38

ROSENZWEIG, MARK S., AND NANDA, NAVIN C. Two-dimensional echocardiographic detection of circulating right atrial thrombi, 435 (Brief comm.)
ROSNER, FRED. (See Zoneraich and Rosner), 927

Ross, David L. (See Mason et al.), 633 Rossi, Peter, Massumi, Ali, Gillette, Paul, and Hall, Robert J. Arrhythmogenic right ventricular dysplasia: Clinical features, diagnostic techniques, and current management. 415

ROULEAU, JEAN-LUCIEN. (See Parmley, Rouleau, and Chatterjee),

ROWEN, MARSHALL. (See Singer, Rowen, and Dorsey), 131

RUBIN, DAVID. (See Kay, Blake, and Rubin), 308 RUBIN, RICHARD E. (See Waller et al.), 1071 RUBIN, STANLEY A. (See Schiff et al.), 438 — (See Swan, Shah, and Rubin), 707

RUDE, ROBERT E. (See Nicod et al.), 822

-, GLOGAR, DIETMAR, KHURI, SHUKRI F., KONER, ROBERT A., GLOGAR, DIETMAR, KHURI, SHUKRI F., KONER, ROBERT A., KARAFFA, STEPHANIE, MULLER, JAMES E., CLARK, LELAND C., JR., AND BRAUNWALD, EUGENE. Effects of intravenous fluorocarbons during and without oxygen enhancement on acute myocardial ischemic injury assessed by measurement of intramyocardial gas tensions, 986

Runco, Vincent. (See Nair et al.), 226 Russell, Richard O., Jr. (See Rackley et al.), 154

RUTITZKY, BERNARDO, GIROTTI, ARNOLDO L., AND ROSENBAUM, MAURICIO B. Efficacy of chronic amiodarone therapy in patients with variant angina pectoris and inhibition of ergonovine coronary constriction, 38 Rutkowski, Monika. (See Valdes-Cruz et al.), 505

SACHS, IRENE. (See Devereux and Sachs), 1086 SAFAR, M. E. (See Payen et al.), 379 SAHN, DAVID J. (See Friedman et al.), 194 SARN, DAVID . (See Valdes-Cruz et al.), 505

— (See Valdes-Cruz et al.), 505

— (See Goldberg et al.), 858
SARTO, TOMOAKI. (See Gettes et al.), 667
SAKSENA, SANJEEV. (See Kessler et al.), 139 SALCEDO, ERNESTO E. (See Mellino et al.), 133

SALCEDO, ERNESTO E. (See Mellino et al.), 222

— (See Chaignon et al.), 374

SALINAS, MATIAS. (See Moncada et al.), 263

SAMLOWSKI, WOLFRAM. (See Wizenberg et al.), 468 Sanderson, Tamara L. (See Kessler et al.), 139 Sands, Militon J. (See Lewis et al.), 498 Scagliotti, Daniel. (See Strasberg et al.), 795

—. (See Bauernfeind et al.), 941 SCHELBERT, HEINRICH R. (See Schön et al.), 532

(See Schön et al.), 548 HENZE, EBERHARD, PHELPS, MICHAEL E., AND KUHL, DAVID E. Assessment of regional myocardial ischemia by posi-

tron-emission computed tomography, 588
Schiff, Steven, Moffatt, Roy, Mandel, William, J., and Rubin,
Stanley A. Acute myocardial infarction and recurrent ventricular arrhythmias in Behcet's syndrome, 438 (Brief comm.)

-, BATEMAN, TIM, MOFFATT, ROY, DAVIDSON, ROBERT, AND BERMAN, DANIEL. Diagnostic considerations in cardiomyopathy: Unique scintigraphic pattern of diffuse biventricular technetium-99m-pyrophosphate uptake in amyloid heart disease, 562 (Brief comm.) Schifferli, J. A. Primary renal origin of hematuria: Importance

of RBC casts and urinary sediment exam technique, 573 (Letter to Editor)

Schmidt, Donald. (See Pollock et al.), 363
Schon, Hans R., Schelbert, Heinrich R., Robinson, Gerald,
Najafi, Ahmad, Huang, Sung-Cheng, Hansen, HerBert, Barrio, Jorge, Kuhl, David E., and Phelps,
Michael E. C-11 labeled palmitic acid for the noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography I. Kinetics of C-11 palmitic acid in normal myocardium,

 Schelbert, Heinrich R., Najafi, Ahmad, Hansen, Herbert, Huang, Henry, Barrio, Jorge, and Phelps, Michael E. C-11 labeled palmitic acid for the noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography II. Kinetics of C-11 palmitic acid in acutely ischemic myocardium, 548

SCHROEDER, JOHN S. (See Ginsburg et al.), 44

. (See Mitchell et al.), 57
 Shah, Pravin M. (See Tei, Shah, and Tanaka), 294

— (See Tei, Shah, and Ormistron), 1025 — (See Cherian et al.), 1074 SHAH, PREDIMAN K. (See Swan, Shah, and Rubin), 707 SHARMA, BIM. (See Francis, Sharma, and Hodges), 995

SHAVIT, SARA. (See Korczyn et al.), 443

SHEEHAN, FLORENCE H., AND EPSTEIN, STEPHEN E. Effects of calcium channel blocking agents on reperfusion

arrhythmias, 973
SHEN, EDWARD Ñ., AND SUNG, RUEY J. Initiation of atrial fibrillation by spontaneous ventricular premature beats in concealed Wolff-Parkinson-White syndrome, 911 (Brief comm.)

Sheps, D. S., Petrovick, M. L., Kizakevich, P. N., Wolfe, C., and Craige, E. Continuous noninvasive monitoring of left ventricular function during exercise by thoracic impedance cardiography-automated derivation of systolic time intervals, 519

SHIH, JOUNG-YI. (See Karpawich et al.), 1001

Shimokawa, Hiroaki, Marsuguchi, Takeyuki, Koiwaya, Yasushi, Fukuyama, Takaya, Orita, Yashuhiko, and Nakamura, Motoomi. Variable exercise capacity in variant angina and greater exertional thallium-201 myocardial defect during vasospastic ischemic ST segment elevation than with ST depression, 142 (Brief comm.)

SIEGEL, ROBERT J., AND ROBERTS, WILLIAM C. Electrocardiographic observations in severe acrtic valve stenosis: Correla-

tive necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to peak systolic transaortic pressure gradi-

ent, 210
Simon, Toby L. (See Bell and Simon), 239
Singer, Melville I., Rowen, Marshall, and Dorsey, Thomas J.
Transluminal acrtic balloon angioplasty for coarctation of the aorta in the newborn, 131 (Brief comm.) Singh, Bramah N. (See Nademanee et al.), 802

—. (See Nademanee et al.), 950
—. (See Josephson et al.), 978
SINGH, R. B., AND CAMERON, E. A. Relation of myocardial magnesium deficiency to sudden death in ischemic heart disease, 449 (Letter to Editor)
SKETCH, MICHAEL H. (See Nair et al.), 226
SKORTON, DAVID J. (See Geiser, Skorton, and Conetta), 905

SLEIGHT, PETER. (See Ramsdale et al.), 459

SLEIGHT, PETER. (See Ramsdale et al.), 459
SMITH, JOAN L. (See Dressendorfer et al.), 358
SMITH, L. RICHARD (See Rogers et al.), 185
SMITH, THOMAS W., BARRY, WILLIAM H., MARSH, JAMES D., AND LORELL, BEVERLY. Hypoxia, calcium fluxes, and inotropic state: Studies in cultured heart cells, 716
SMUCKLER, ALAN L., NOMEIR, ABDEL-MOHSEN, WATTS, L. EARL, AND HACKSHAW, BARRY T. Echocardiographic diagnosis of aortic root dissection by M-mode and two-dimensional techniques. 897 al techniques, 897

Sobel, Burton E. Diagnostic promise of positron tomography, 673

Sobol, Samuel M., Brown, Jerry M., Bunker, Stephen R., Patel, Juibhoy, and Lull, Robert J. Noninvasive diagnosis of cardiac amyloidosis by technetium-99m-pyrophosphate myocardial scintigraphy, 563 (Brief comm.)

SOHN, YOUNG H. (See Wizenberg et al.), 468

Sonn, Jagmeet S. (See Ptacin et al.), 54 Sondheimer, Henry M. (See Kavey, Blackman, and Sondheimer), 342

SONES, F. MASON, Jr. (See Kramer et al.), 283 SONNENBLICK, EDMUND H. (See Kirk and Sonnenblick), 756

— (See Fisher, Sonnenblick, and Kirk), 966 Soumelas, N. (See Venetis et al.), 924 Spain, David M. (See Kossowsky, Lyon, and Spain), 926 Spielman, Scott R. (See Horowitz et al.), 604

SPODICK, DAVID H. (See Doi et al.), 301 STACK, RICHARD S. (See Bardy et al.), 1068 STANG, JOHN M. (See Lewis et al.), 123 STRASBERG, BORIS (See Swiryn et al.), 168 —. (See Swiryn et al.), 319

-, Lam, Wilfred, Swiryn, Steven, Bauernfeind, Robert, Sca-GLIOTTI, DANIEL, PALILEO, EDWIN, AND ROSEN, KENNETH. Symptomatic spontaneous paroxysmal AV nodal block due to localized hyperresponsiveness of the AV

node to vagotonic reflexes, 795

—. (See Bauernfeind et al.), 941
STRAUSS, H. WILLIAM. (See Brown et al.), 1019 STRONG, WILLIAM B. (See Covitz et al.), 1072

Sugura, Tetsuro. (See Doi et al.), 301 Sung, Ruey J. (See Shen and Sung), 911 Surawicz, Borys. Role of calcium-blocking agents in treatment of cardiac arrhythmias related to myocardial ischemia,

SWAN, H. J. C., SHAH, PREDIMAN K., AND RUBIN, STANLEY. Role of vasodilators in the changing phases of acute myocardi-

al infarction, 707
SWERDLOW, CHARLES D. (See Mason et al.), 633
SWIRYN, STEVEN, BAUERNFEIND, ROBERT A., PALILEO, EDWIN A.,
STRASBERG, BORIS, DUFFY, C. ELISE, AND ROSEN, KEN-NETH M. Electrophysiologic study demonstrating triple antegrade AV nodal pathways in patients with spontaneous and/or induced supraventricular tachycardia, 168

—, Pavel, Dan, Byrom, Ernest, Bauernfeind, Robert A., Strasberg, Boris, Palileo, Edwin, Lam, Wilfred, Wyndham, Christopher R. C., and Rosen, Kenneth M. Sequential regional phase mapping of radionuclide gated biventriculograms in patients with sustained ventricular tachycardia: Close correlation with elec-

trophysiologic characteristics, 319
-. (See Strasberg et al.), 795 —. (See Bauernfeind et al.), 941 SYMBAS, P. N. (See Wilcox et al.), 148

TAJIMI, TSUKASA. (See Kaseda et al.), 569 TAKAHASHI, OSAHIR². (See Masuda et al.), 230
TAKSHITA, AKIRA. See Kaseda et al.), 569
TANAKA, HIROMITSU. (See Tei, Shah, and Tanaka), 294
TARAZI, ROBERT C. (See Chaignon et al.), 374

TATENO, YOKIO. (See Masuda et al.), 230
TEI, CHUWA, SHAH, PRAVIN M., AND TANAKA, HIROMITSU. Phonographic-echographic documentation of systolic honk in tricuspid prolapse, 294 (Brief comm.)

-, AND ORMISTON, JOHN A. Assessment of tricuspid regurgitation by directional analysis of right atrial systolic linear reflux echoes with contrast M-mode echocardiography, 1025

(See Cherian et al.), 1074

Testa, R. (See Biagini et al.), 13
Thien, Theo A. (See Huysmans, Thien, and Koene), 395
Thompson, Melish A. (See Olinger, Thompson, and Keelan),

440 THOMSON, A. (See James and Thomson), 457
THRALL, JAMES H. (See Rizi et al.), 292
—. (See Pitt and Thrall), 784
TIEMANN, JEROME J. (See Friedman et al.), 194
TOKUNAGA, KOUCHL. (See Kaseda et al.), 569
TOTOMOKOUO, J. A. (See Payen et al.), 379
TRESCH, DONALD D. (See Ptacin et al.), 54
TRESUS, SALVADOR, (See Harritz et al.), 421

TREVES, SALVADOR. (See Hurwitz et al.), 421 TRONCOSO, PATRICIA. (See Mehlman et al.), 145

UCHIYAMA, Gyo. (See Masuda et al.), 230

ULMSTEN, ULF, BERNSTEIN, KURT, AND MATTSON, SÖREN. The nique for localization and dynamic evaluation of limb deep vein thrombosis, 448 (Letter to Editor)

UNTEREKER, WILLIAM J. (See Martin et al.), 960

VALDES-CRUZ, LILLIAM M., REINECKE, CYNTHIA, RUTKOWSKI, MONI-KA, DUDELL, GOLDE G., GOLDBERG, STANLEY J., ALLEN, HUGH D., SAHN, DAVID J., AND PIOMELLI, SERGIO. Pre-clinical abnormal segmental cardiac manifestations of thalassemia major in children on transfusion-chelation therapy: Echographic alterations of left ventricular posterior wall contraction and relaxation patterns, 505

—. (See Goldberg et al.), 858 VALENSTEIN, PAUL. (See Bardy et al.), 1068 VALOVIS, RICARDO, AND CHARUZI, YZHAR. Vegetation diastolic sound in mitral valve endocarditis, 432 (Brief comm.)

VANDENBOSSCHE, JEAN-LUC, AND ENGLERT, MARC. Effects of respiration on an atrial septal aneurysm of the fossa ovale shown by echographic study, 922 (Brief comm.) VASILEIADES, J. (See Venetis et al.), 924

VASILEIADES, J. (See Venetis et al.), 924
VASUDEVAN, GOPALAN. (See Mellino et al.), 222
VENETIS, C., SOUMELAS, N., NIHOYANNOPOULOS, P., AND VASILEIADES,
J. Shortened aortic ejection period with consequent
apparent early diastolic murmur in mitral valve prolapse syndrome, 924 (Brief comm.)
VERA, ZAKAUDDIN, AWAN, NAJAM A., AND MASON, DEAN T. Assess-

ment of oral quinidine effects on sinus node function

in sick sinus syndrome patients, 80 Vidne, Bernardo. (See Korczyn et al.), 443

Wackers, Frans J. Th. (See Lewis et al.), 498
Waeber, Bernard, Gavras, Irene, Brunner, Hans R., Cook,
Charles A., Charocopos, Fotis, and Gavras, Haralamnos P. Prediction of sustained antihypertensive effica-cy of chronic captopril therapy: Relationships to immediate blood pressure response and control plas-

ma renin activity, 384 Wagner, Henry N., Jr. Nuclear techniques in ischemic heart disease, 681

WALINSKY, PAUL. Angiographic documentation of spontaneous spasm of saphenous vein coronary artery bypass graft, 290 (Brief comm.)

WALLER, BRUCE F. (See McManus et al.), 451

—, RUBIN, RICHARD E., McGrath, Francis J., Del Negro, Albert
A., AND ROBERTS, WILLIAM C. Coronary calcium—a clue to angiographic underestimation of coronary luminal narrowing, 1071 (Brief comm.)
WALLIS, DIANE. (See O'Connell et al.), 85

WARD, ANN. (See Pollock et al.), 363

WARD, CHRISTOPHER. (See Ramsdale et al.), 459
WARREN, JAMES V. (See Lewis et al.), 123
WATANABE, AUGUST M. (See Lindemann, Bailey, and Watanabe),

WATANABE, SHIGERU. (See Masuda et al.), 230

Watkins, Levi. (See Mirowski et al.), 147 Watts, L. Earl. (See Smuckler et al.), 897 Weber, Karl T., Andrews, Virginia, Janicki, Joseph S. Cardiotonic agents in the management of chronic cardiac

tonic agents in the management of chronic cardiac failure, 639
Weiss, Y. A. (See Payen et al.), 379
Weissler, Arnold M. (See Wizenberg et al.), 468
Weld, Francis M. (See Bigger, Weld, and Roinitzky), 660
Wellens, Hein J. J. (See de Zwaan, Bär, and Wellens), 730
West, C. E. (See Knuiman, West, and Hautvast), 447
West, R. R. Early mobilization after uncomplicated myocardial infarction, 311 (Letter to Editor)
Whittle, James L., Feldman, Robert L., Pepine, Carl J., Curry, R. Charles, and Conti, C. Richard. Variability of electrocardiographic responses to repeated ergonovine provocation in variant angina patients with coronary provocation in variant angina patients with coronary

provocation in variant angina patients with coronary artery spasm, 161
Wiener, Isaac. (See Kay, Estioko, and Wiener), 338
Wikman-Coffelt, Joan. (See Riemenschneider et al.), 834
—, Parmley, William W., and Mason, Dean T. Relation of myosin isozymes to the heart as a pump, 934
Wilcox, W. Dean, Plauth, W. H., Jr., Williams, Willis H., and Symbas, P. N. Immediate postoperative regurgitant malfunction of Björk-Shiley aortic valve due to interfering Teflon ventricular septal defect repair: Correction by rotation of prosthetic annulus, 148 (Brief tion by rotation of prosthetic anulus, 148 (Brief comm.)

COMM.)
WILLERSON, JAMES T. (See Johnson et al.), 4
—. (See Nicod et al.), 822
WILLIAMS, WILLIS H. (See Wilcox et al.), 148
WINKLE, ROGER A. (See Bragg-Remschel, Anderson, and Winkle),

—. (See Keefe, Peters, and Winkle), 511 —. (See Mason et al.), 633 Winniford, Michael D. (See Johnson et al.), 4 Wizenberg, Theodore A., Muz, Jaroslaw, Sohn, Young H., Samlowski, Wolfram, and Weissler, Arnold M. Value of positive myocardial technetium-99m-pyrophosphate scintigraphy in the noninvasive diagnosis of cardiac amyloidosis, 468
Wolfe, C. (See Sheps et al.), 519
Wong, Maylene. (See Cherian et al.), 1074

Wood, Earl H., Gilbert, Barry K., Harris, Lowell D., Kinsey, James H., Ritman, Erik L., and Robb, Richard A. High temporal resolution synchronous volumetric scanning tomography: Potential roles in clinical evaluation of

woosley, Raymond L., Roden, Dan M., Duff, Henry J., and Oates, John A. Selection of an antiarrhythmic drug for a sudden-death-prevention trial, 737

WYNDHAM, CHRISTOPHER R. C. (See Swiryn et al.), 319

YAGIL, Y., KOBRIN, I., LEIBEL, B., AND BEN-ISHAY, D. Ischemic ECG changes with initial nifedipine therapy of severe hypertension, 310 (Letter to Editor)

Yoshida, Hideo. (See Masuda et al.), 230 YUSUF, SALIM. (See Ramsdale et al.), 459

ZAMAN, LIAQUAT. (See Myerburg et al.), 615 ZARET, BARRY L. (See Lewis et al.), 498 ZELINGER, ALLAN B., FALK, RODNEY H., AND HOOD, WILLIAM B., JR. Electrical-induced sustained myocardial depolariza-tion as a possible cause for transient ST elevation post-DC elective cardioversion, 1073 (Brief comm.)

ZIBELLI, LOU'S R. (See Hecht and Zibelli), 436
ZINNER, ALEX. (See Karpawich et al.), 1001
ZIPES, DOUGLAS P. (See Rinkenberger et al.), 177
— (See Prystowsky et al.), 426
—, PRYSTOWSKY, E. N., AND HEGER, J. J. Electrophysiologic testing

of antiarrhythmic drug agents, 610

Jackman, Warren M., Heger, James J., Chilson, Donald A.,
Browne, Keven F., Naccarelli, Gerald V., Rahilly,
G. Thomas, J., and Physrowsky, Eric N. Clinical
transvenous cardioversion of recurrent life-threatentransvenous cardioversion of recurrent life-threaten-ing ventricular tachyarrhythmias: Low energy syn-chronized cardioversion of ventricular tachycardia and termination of ventricular fibrillation in patients using a catheter electrode, 789 Zoneraich, Samuel, and Rosner, Fred. Third comment, 927 (Letter to Editor)

Subject index*

Accessory connections, right anterior, prolonged and decremental antegrade conduction properties in: wide QRS anti-dromic tachycardia of left bundle branch block pat-tern without Wolff-Parkinson-White configuration in

sinus rhythm (Gillette et al.), 66
pathway, concealed, post-myocardial infarction incessant
supraventricular tachycardia due to (Prystowsky et al.), 426 (Brief comm.)

Acute allergic reaction with coronary vasospasm (Hirsh), 928 (Letter)

Reply (Ginsburg and Bristow), 928

Adenosine triphosphatase activity, cardiac muscle myosin, relative to hemodynamic alterations in newborn lambs, maturational changes in (Riemenschneider et al.),

Adenovirus infection and Balkan nephritis (Georgescu et al.), 572 (Letter)

Allergic reaction, acute, with coronary vasospasm (Hirsh), 928 (Letter) Reply (Ginsburg and Bristow), 928

*January, pp. 1-160; February, pp. 161-318; March, pp. 319-458; April, Part 1, pp. 459-582; April, Part 2, pp. 583-788; May, pp. 789-940; June, pp. 941-1128.

Ambulatory ECG monitoring systems, frequency response characteristics of and their implications for ST segment analysis (Bragg-Remschel, Anderson, Winkle), 20

management of severe chronic congestive heart failure, long-term hemodynamic and clinical efficacy of captopril therapy in (Awan et al.), 474

Amines, sympathomimetic: potential clinical applications in is-

Amines, sympathonimete: botendar clinical applications in schemic heart disease (Goldberg and Raifer), 724
Aminophylline therapy and mechanisms: paroxysmal nocturnal dyspnea in CHF (Rees and Clark), 311 (Letter)
Amiodarone in patients with life-threatening ventricular arrhythmias, antiarrhythmic efficacy and electrophysiologic actions of: potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of induced ventricular tachycardia (Nademanee et al.),

therapy, chronic, efficacy of in patients with variant angina pectoris and inhibition of ergonovine coronary constriction (Rutitzky, Girotti, and Rosenbaum), 38

Amyloid heart disease, unique scintigraphic pattern of diffuse biventricular technetium-99m-pyrophosphate uptake in: diagnostic considerations in cardiomyopathy (Schiff et al.), 562 (Brief comm.)

Amyloidosis, cardiac, noninvasive diagnosis of by technetium-99m-pyrophosphate myocardial scintigraphy (Sobol et al.), 563 (Brief comm.)

- Amyloidosis, cardiac-continued
 - value of positive myocardial technetium-99m-pyrophosphate scintigraphy in the noninvasive diagnosis of (Wizenberg et al.), 468
- Analog method, accurate rapid compact, for quantification of frequency and duration of myocardial ischemia by semiautomated analysis of 24-hour Holter ECG recordings (Nademanee et al.), 802
- Aneurysm, atrial septal, of the fossa ovale, effects of respiration on an shown by echographic study (Vandenbossche and Englert), 922 (Brief comm.)
- huge false, due to rupture of right coronary artery in Behcet's syndrome (Kaseda et al.), 569 (Brief comm.)
- submitral left ventricular, two-dimensional echocardiographic
- features of (Davis et al.), 289 (Brief comm.)

 Aneurysmectomy and endocardial resection for ventricular tachycardia: favorable hemodynamic and antiarrhythmic results in patients with global left ventricular dysfunction (Martin et al.), 960
- Angina patients, unstable, vasospastic ischemic mechanism of frequent asymptomatic transient ST-T changes during continuous electrocardiographic monitoring in selected (Biagini et al.), 13
 - variant, with coronary artery spasm, variability of electrocardiographic responses to repeated ergonovine provocation in (Whittle et al.), 161
- pectoris and myocardial ischemia, role of coronary vasospasm in the pathogenesis of (Gorlin), 598 stable, coronary patients with, effects of beta-adrenergic blockade with pindolol versus placebo in (Dwyer, Pepe, and Pinkernell), 830
 - unstable, continuous electrocardiographic monitoring in patients with: identification of high-risk subgroup with severe coronary disease, variant angina, and/or
 - impaired early prognosis (Johnson et al.), 4 : is it time to change our approach? (Rackley et al.), 154
 - variant, and inhibition of ergonovine coronary constriction, efficacy of chronic amiodarone therapy in patients with (Rutitzky, Girotti, and Rosenbaum), 38
- due to coronary artery spasm, randomized double-blind comparison of nifedipine and isosorbide dinitrate therapy in (Ginsburg et ɛ!), 44 phenomenon, walk-through, demonstrated by graded exercise radionuclide ventriculography: possible coronary spasm mechanisms (Rizi et al.), 292 (Brief comm.)
- variant, severe coronary disease, and/or impaired early prognosis, identification of high-risk subgroup with: continuous electrocardiographic monitoring in patients with unstable angina pectoris (Johnson et al.), 4
- variable exercise capacity in and greater exertional thallium-201 myocardial defect during vasospastic ischemic ST segment elevation than with ST depression (Shimokawa et al.), 142 (Brief comm.)
 vasospastic, ergonovine-induced, negative U waves and peaked T waves without ST changes during spontaneous and
- (Matsuguchi et al.), 918 (Brief comm.)
- Angiocardiographic, electrocardiographic, and echocardiographic pre- and postoperative evaluation of primary benign intramural ventricular tumors in children (Biancaniel-
- lo et al.), 852
 Angiocardiography, radionuclide, current value of for shunt quantification and management in patients with ecundum atrial septal defect (Hurwitz et al.), 421
- Angiographic documentation of spontaneous spasm of saphenous vein coronary artery bypass graft (Walinsky), 290 (Brief comm.)
- underestimation of coronary luminal narrowing-coronary calcium (Waller et al.), 1071 (Brief comm.)
- Angiography, coronary, during acute myocardial infarction: a search for spasm (Leinbach and Gold), 768 radionuclide, first-pass, in visualization of acute myocardial infarction ventricular septal defect and assessment of ventricular function (Hecht and Zibelli), 436 (Brief
- Angioplasty, coronary, and implications for the future, results
 - from (Gruentzig), 779
 percutaneous transluminal (PTCA) and percutaneous transluminal coronary recanalization (PTCR), successful treatment of acute myocardial infarction shock by combined (Meyer et al.), 132 (Brief comm.)

- Angioplasty-continued
 - transluminal aortic balloon, for coarctation of the aorta in the newborn (Singer, Rowen, and Dorsey), 131 (Brief
- Anomalous pulmonary venous connection, total: survival for 62
- years without surgical intervention (McManus, Luetzeler, and Roberts), 298 (Brief comm.)
 Antiarrhythmic drug agents, electrophysiologic testing of (Zipes, Prystowsky, and Heger), 610
 - selection of an for a sudden-death-prevention trial (Woosley
 - et al.), 737
 efficacy and electrophysiologic actions of amiodarone in patients with life-threatening ventricular arrhythmias: potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of induced ventricular tachycardia (Nademanee et al.), 950
- therapy, programmed ventricular stimulation in predicting vulnerability to ventricular arrhythmias and their response to (Mason et al.), 633
- Anticoagulation, long-term oral coumarin, prediction of hemorrhage during by excessive prothrombin ratio (Forfar), 445 (Brief comm.)
- Antidepressant drugs, tricyclic, cardiovascular effects of: therapeutic usage, overdose, and management of complications (Marshall and Forker), 401
- Antihypertensive efficacy of chronic captopril therapy, prediction of sustained: relationships to immediate blood pressure response and control plasma renin activity (Waeber et al.), 384
- Antihyperviscosity effect of hemodilution with stroma-free hemoglobin, early deleterious hemorheologic changes following acute experimental coronary occlusion and salutary (Biro, Beresford-Kroeger, and Hendry),
- Antilog signal processing of two-dimensional echocardiographic images, high predictive accuracy for detection of left main coronary artery disease by (Friedman et al.), 194
- Anulus calcification, mitral, echographic-quantified severity of: prognostic correlation to related hemodynamic, valvular, rhythm, and conduction abnormalities (Mellino et
- al.), 222 prosthetic, correction by rotation of: immediate postoperative regurgitant malfunction of Björk-Shiley aortic valve due to interfering Teflon ventricular septal defect
- repair (Wilcox et al.), 148 (Brief comm.) Aorta in the newborn, coarctation of the, transluminal aortic balloon angioplasty for (Singer, Rowen, and Dorsey), 131 (Brief comm.)
- Aortic balloon angioplasty, transluminal, for coarctation of the aorta in the newborn (Singer, Rowen, and Dorsey), 131 (Brief comm.)
- dissection in Turner's syndrome (Lie), 1077 (Brief comm.) ejection period, shortened, with consequent apparent early diastolic murmur in mitral valve prolapse syndrome
- (Venetis et al.), 924 (Brief comm.)
 pressure, central, noninvasive estimation of using the oscillometric method for analyzing systemic artery pulsatile blood flow: comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure mea
 - surements (Borow and Newburger), 879 measurements, comparative study of indirect systolic, dia stolic, and mean brachial artery pressure with simultaneous direct ascending: noninvasive estimation of central aortic pressure using the oscillometric method for analyzing systemic artery pulsatile blood flow (Borow and Newburger), 879
- regurgitation, chronic, radionuclide angiographic exercise LV performance in: relationship to resting echographic ventricular dimensions and systolic wall stress index (Lewis et al.), 498
- utility of two-dimensional echocardiography in the differential diagnosis of the etiology of (Imaizumi et al.),
- root dissection, echocardiographic diagnosis of by M-mode and two-dimensional techniques (Smuckler et al.), 897 stenosis, valvular, range gated echo-Doppler velocity and tur-
- bulence mapping in patients with (Goldberg et al.),

Aortic stenosis, valvular-continued

severe, identification of in adult patients: reliable estima-tion of peak left ventricular systolic pressure by M-mode echographic-determined end-diastolic relative wall thickness (Reichek and Devereux), 202

valve disease patients with or without concomitant coronary disease, prevalence and signficance of reversible radio-nuclide ischemic perfusion defects in symptomatic

(Pfisterer et al.), 92

prosthesis, competent, downward displacement of without left ventricular enlargement: a new sign of ascending root dissection in absence of valvular regurgitation (Gooch), 920 (Brief comm.)

stenosis, severe, electrocardiographic observations in: correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to peak systolic transaortic pressure gradient (Siegel and Roberts), 210

Aortocaval fistula-induced chronic left ventricular volume overload and normal systolic performance, comparative hemodynamic effects of digoxin vs nitroprusside in conscious dogs with (O'Rourke, Badke, and Forst), 489

Approach to management of unstable angina pectoris-is it time to change? (Rackley et al.), 154

Arrest, cardiac, in pheochromocytoma (Ouzts and Newland), 928 (Letter)

Arrhythmia(s), cardiac, related to myocardial ischemia, role of calcium-blocking agents in treatment of (Surawicz), 698

life-threatening, out-of-hospital, successful conversion of with the implanted automatic defibrillator (Mirowski et al.), 147 (Brief comm.)

postinfarction ventricular—which should be treated? (Bigger, Weld, and Rolnitzky), 660
reperfusion, effects of calcium channel blocking agents on (Sheehan and Epstein), 973

stable and potentially lethal, evolving concepts of management

of (Myerburg et al.), 615 ventricular, and their response to antiarrhythmic therapy, programmed ventricular stimulation in predicting vulnerability to (Mason et al.), 633

life-threatening, antiarrhythmic efficacy and electrophysiologic actions of amiodarone in patients with: potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of induced ventricular tachycardia (Nademanee et al.), 950

numerators and denominators revisited relative to electrophysiologic study of: the stampede to stimulation (Graboys), 1089

recurrent, and acute myocardial infarction in Behcet's syndrome (Schiff et al.), 438 (Brief comm.)

role of programmed stimulation in assessing vulnerability to (Horowitz et al.), 604

vulnerability in acute myocardial infarction, factors related to (Gettes et al.), 667

Arrhythmogenic right ventricular dysplasia: clinical features, diagnostic techniques, and current management (Rossi et al.), 415 two-dimensional echocardiographic detection of (Baran

et al.), 1066 (Brief comm.) Arrhythmogenicity, altered, catecholamine-enhanced positive chronotropy without: denervation supersensitivity in the canine heart (Korczyn et al.), 443 (Brief comm.)

Arterial disorders, peripheral, chronic, beneficial hemorheologic therapy of with pentoxifylline: results of double-blind study versus vasodilator-nylidrin (Accetto), 864

Artery disease, coronary, heart failure secondary to, vasodilators in (Parmley, Rouleau, and Chatterjee), 625

left anterior descending, benign and frequent condition involving the: clinical significance of isolated coronary bridges (Kramer et al.), 283

spasm, coronary: prevalence, clinical significance, and provoca-tive testing (Conti, Feldman, and Pepine), 584 randomized double-blind comparison of nifedipine and isosorbide dinitrate therapy in variant angina pectoris

due to (Ginsburg et al.), 44 ASH in conditions other than hypertrophic cardiomyopathy (Ramsdale and Bennett), 1083 (Letter) Ataxia, Friedreich's, characteristics of the cardiac hypertrophy in

(Gottdiener et al.), 525

Atenolol, intravenous, ischemic pain relief in patients with acute myocardial infarction by (Ramsdale et al.), 459

Atherosclerosis, coronary, and acute myocardial ischemic attacks,

Atheroscierosis, coronary, and acute myocardial ischemic attacks, role of hypertension and downward changes of blood pressure in the genesis of (Green), 579

Atresia, tricuspid, with previous Glenn shunt, assessment of Fontan graft patency by radionuclide perfusion pulmonary scan in (Covitz et al.), 1072 (Brief comm.)

Atrial fibrillation, initiation of by spontaneous ventricular premature beats in concealed Wolff-Parkinson-White syndrome (Shen and Sung), 911 (Brief comm.)

paroxysmal, sporadic, electrophysiologic drug testing in prophylaxis of: technique, application, and efficacy in everely symptomatic preexcitation patients (Bauernfeind et al.), 941

rhythm, left, clinical confirmation of ECG criteria for (Beder et al.), 848

septal aneurysm of the fossa ovale, effects of respiration on an shown by echographic study (Vandenbossche and Englert), 922 (Brief comm.)

defect, secundum, current value of radionuclide angiocardiography for shunt quantification and management in patients with (Hurwitz et al.), 421

Atrioventricular (AV) nodal block, paroxysmal, symptomatic spontaneous, due to localized hyperresponsiveness of the AV node to vagotonic reflexes (Strasberg et al.), 795

pathways, triple antegrade, in patients with spontaneous and/or induced supraventricular tachycardia, electrophysiologic study demonstrating (Swiryn et al.), 168

node-fascicular and sinus node-atrial disease in patients with mitral anular calcification, high prevalence of symptomatic bradyarrhythmias due to (Nair et al.), 226 hyperresponsiveness of to vagotonic reflexes, symptomatic spontaneous paroxysmal AV nodal block due to local

ized (Strasberg et al.), 795

pacing, sequential, in right ventricular myocardial infarction with complete heart block (Abraham and Norris), 929 (Letter)

Axial oblique contrast left ventriculography, quantitative: valida-tion of the method by demonstrating improved visual-ization of regional wall motion and mitral valve function with accurate volume determinations (Rogers et al.), 185

Balkan nephritis and adenovirus infection (Georgescu et al.), 572 (Letter)

Balloon angioplasty, aortic, transluminal, for coarctation of the aorta in the newborn (Singer, Rowen, and Dorsey), 131 (Brief comm.)

Behcet's disease, recognition of the diverse cardiovascular manifestations in (James and Thomson), 457
syndrome, acute myocardial infarction and recurrent ventricular arrhythmias in (Schiff et al.), 438 (Brief comm.)
huge false aneurysm due to rupture of the right coronary

artery in (Kaseda et al.), 569 (Brief comm.) Benign intramural ventricular tumors in children, primary: pre-and postoperative ECG, echocardiographic, and angiocardiographic evaluation (Biancaniello et al.),

Beta-adrenergic blockade with pindolol versus placebo in coro-

nary patients with stable angina pectoris, effects of (Dwyer, Pepe, and Pinkernell), 830 blocking agent and diazoxide, combined intravenous administration of in acute treatment of severe hypertension or hypertensive crisis (Huysmans, Thien, and Koene),

Bicuspid pulmonic valve, midsystolic click accompanying isolated (Mehlman et al.), 145 (Brief comm.)

Biochemical mechanisms, potential, for regulation of the slow inward current: theoretical basis for drug action (Lin-

demann, Bailey, and Watanabe), 746 Biventricular technetium-99m-pyrophosphate uptake in amyloid heart disease, unique scintigraphic pattern of diffuse: diagnostic considerations in cardiomyopathy (Schiff et al.), 562 (Brief comm.)

- Biventriculograms, radionuclide gated, sequential regional phase mapping of in patients with sustained ventricular tachycardia: close correlation with electrophysiologic characteristics (Swiryn et al.), 319
- Björk-Shiley aortic valve, immediate postoperative regurgitant malfunction of due to interfering Teflon ventricular
 - septal defect repair: correction by rotation of pros-thetic anulus (Wilcox et al.), 148 (Brief comm.) unmarked tilting disc mitral valve prosthesis, optimal man-agement of suspected thrombosis of standard (Olinger, Thompson, and Keelan), 440 (Brief comm.)
- Blocking agents, calcium channel, effects of on reperfusion arrhythmias (Sheehan and Epstein), 973 Blood pressure, downward changes of, and hypertension, role of in the genesis of coronary atherosclerosis and acute
 - myocardial ischemic attacks (Green), 579
 - independent effects of Na reduction and K supplementation
 - on (Gillum, Prineas, and Kebede), 449 (Letter) response and control plasma renin activity, relationships to immediate: prediction of sustained antihypertensive efficacy of chronic captopril therapy (Waeber et al.),
 - tilt-test response, increased diastolic orthostatic, prognostic value of and subsequent weight gain: prospective study of predictive factors determining borderline hypertensive individuals who develop sustained hypertension (Payen et al.), 379
- Bookshelf, 160, 1096 Borderline hypertensive individuals who develop sustained hypertension, prospective study of predictive factors determining: prognostic value of increased diastolic orthostatic blood pressure tilt-test response and subsequent weight gain (Payen et al.), 379
- Brachial artery pressure, mean, comparative study of indirect systolic, diastolic, and with simultaneous direct ascending aortic pressure measurements: noninvasive estimation of central aortic pressure using the oscillometric method for analyzing systemic artery pulsatile blood flow (Borow and Newburger), 879
- Bradyarrhythmias, symptomatic, high prevalence of due to atrioventricular node-fascicular and sinus node-atrial disease in patients with mitral anular calcification (Nair et al.), 226
- Bridges, coronary, isolated, clinical significance of: benign and frequent condition involving the left anterior descending artery (Kramer et al.), 283
- Bundle branch block, transient, following use of hypothermic cardioplegia in coronary artery bypass surgery: high incidence without perioperative myocardial infarction (O'Connell et al.), 85
- Bypass graft, saphenous vein coronary artery, angiographic docu-mentation of spontaneous spasm of (Walinsky), 290 (Brief comm.)
 - surgery, coronary artery, transient bundle branch block following use of hypothermic cardioplegia in: high incidence without perioperative myocardial infarction (O'Connell et al.), 85
 - C
- C-11 labeled palmitic acid for noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography. I. Kinetics of C-11 palmitic acid in normal myocardium (Schön et al.), 532

 II. Kinetics of C-11 palmitic acid in acutely ischemic myocardium (Schön et al.), 548

 - palmitic acid in acutely ischemic myocardium, kinetics of. Part II of C-11 labeled palmitic acid for the noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography (Schön et
 - in normal myocardium, kinetics of. Part I of C-11 labeled palmitic acid for the noninvasive evaluation of region-
- al myocardial fatty acid metabolism with positron-computed tomography (Schön et al.), 532 Calcification, mitral anular, high prevalence of symptomatic bradyarrhythmias due to atrioventricular node-fascicular and sinus node-atrial disease in patients with (Nair et al.), 226

- Calcium-blocking agents in treatment of cardiac arrhythmias related to myocardial ischemia, role of (Surawicz), 698
 - channel blocking agents, effects of on reperfusion arrhythmias (Sheehan and Epstein), 973 coronary—a clue to angiographic underestimation of coronary
 - luminal narrowing (Waller et al.), 1071 (Brief comm.)
 - fluxes, hypoxia, and inotropic state: studies in cultured heart cells (Smith et al.), 716
 serum increased, with reduced potassium despite hypovolemic-
 - reduced cardiac output, improved contractility resulting from: acute effects of hemodialysis on echographic-determined cardiac performance (Chaignon et al.).
- Captopril, chronic, and diuretic therapy of severe congestive heart failure, postural hypotension during tilt with (Cody, Franklin, and Laragh), 480
 - therapy, chronic, prediction of sustained antihypertensive effi-cacy of: relationships to immediate blood pressure response and control plasma renin activity (Waeber et
 - for severe CHF: hypotensive response in presence of marked-ly elevated PRA (LaBarre, O'Connell, and Gunnar), 308 (Letter)
 - in ambulatory management of severe chronic congestive heart failure, long-term hemodynamic and clinical efficacy of (Awan et al.), 474
- Cardiac amyloidosis, noninvasive diagnosis of by technetium-99m-pyrophosphate myocardial scintigraphy (Sobol et al.), 563 (Brief comm.)
 - value of positive myocardial technetium-99m-pyrophosphate scintigraphy in the noninvasive diagnosis of (Wizenberg et al.), 468
 - arrest in pheochromocytoma (Ouzts and Newland), 928 (Letter)
 - prehospital, sudden coronary death survivors of, evaluation of postresuscitation left ventricular global and segmental function by radionuclide ventriculography in: correlation to subsequent short-term prognosis (Ptacin et al.), 54
 - arrhythmias related to myocardial ischemia, role of cal-cium-blocking agents in treatment of (Surawicz),
- care, prehospital: does it make a difference? (Cobb), 316 catheterization, no, before cardiac valve replacement-a mistake (Roberts), 930
- failure, chronic, cardiotonic agents in the management of (Weber, Andrews, and Janicki), 639 hypertrophy in Friedreich's ataxia, characteristics of the (Gott-
- diener et al.), 525
- lung, shock lung, and principles of ventilatory therapy in adult respiratory distress syndrome: mechanisms and conse-
- quences of pulmonary edema (Ayres), 97 manifestations of thalassemia major in children on transfusionchelation therapy, preclinical abnormal segmental: echographic alterations of left ventricular posterior wall contraction and relaxation patterns (Valdes-Cruz et al.), 505
- motion, total and regional, ECG synchronized computed tomography in clinical evaluation of: comparison of postmyocardial infarction to normal hearts by rapid sequential imaging (Masuda et al.), 230
- muscle myosin adenosine triphosphatase activity relative to hemodynamic alterations in newborn lambs, maturational changes in (Riemenschneider et al.), 834
- output, hypovolemic-reduced, improved contractility resulting from serum increased calcium with reduced potassium despite: acute effects of hemodialysis on echographicdetermined cardiac performance (Chaignon et al.),
- pain, consideration of the fundamental mechanisms eliciting (Malliani and Lombardi), 575
- performance, acute effects of hemodialysis on echographic-determined: improved contractility resulting from serum increased calcium with reduced potassium despite hypovolemic-reduced cardiac output (Chaignon et al.), 374
- valve replacement, complications of: characteristic abnormalities of prostheses pertaining to any or specific site (Roberts), 113

- Cardiac valve replacement-continued
 - no cardiac catheterization before-a mistake (Roberts), 930
- Cardiocirculatory diseases, feasibility of intravascular laser irra-diation for in vivo visualization and therapy of (Lee et al.), 1076 (Brief comm.)
- al.), 1076 (Brief comm.)

 Cardiogenic circulatory collapse, acute, comparative hemodynamic effects of dopamine and dobutamine in patients with (Francis, Sharma, and Hodges), 995

 Cardiomyopathy, diagnostic considerations in: unique scintigraphic pattern of diffuse biventricular technetium-99m-pyrophosphate uptake in amyloid heart disease (Schiff et al.), 562 (Brief comm.)

 hypertrophic, ASH in conditions other than (Ramsdale and Bennett), 1083 (Letter)

 Cardioplegia, hypothermic, in coronary artery bypass surgery, transient bundle branch block following use of: high incidence without perioperative myocardial infarction (O'Connell et al.), 85
- (O'Connell et al.), 85
- Cardiotonic agents in the management of chronic cardiac failure (Weber, Andrews, and Janicki), 639
- Cardiovascular diagnostic technique, the emergence of computed tomography as a (Brundage and Lipton), 313
- effects of tricyclic antidepressant drugs: therapeutic usage, overdose, and management of complications (Mar-shall and Forker), 401
- manifestations, diverse, in Behcet's disease, recognition of the
- (James and Thomson), 457 Cardioversion, clinical transvenous, of recurrent life-threatening ventricular tachyarrhythmias: low energy synchro-nized cardioversion of ventricular tachycardia and termination of ventricular fibrillation in patients using a catheter electrode (Zipes et al.), 789
- low energy synchronized, of ventricular tachycardia and termination of ventricular fibrillation in patients using a catheter electrode: clinical transvenous cardioversion of recurrent life-threatening ventricular tachyarrhythmias (Zipes et al.), 789
 elective, electrical-induced sustained myocardial
- depolarization as a possible cause for transient ST elevation (Zelinger, Falk, and Hood), 1073 (Brief comm.)
- Catecholamine-enhanced positive chronotropy without altered arrhythmogenicity: denervation supersensitivity in the canine heart (Korczyn et al.), 443 (Brief comm.)
- Catheter electrode, low energy synchronized cardioversion of ventricular tachycardia and termination of ventricular fibrillation in patients using a: clinical transvenous cardioversion of recurrent life-threatening ventricular tachyarrhythmias (Zipes et al.), 789
- Catheterization, cardiac, no, before cardiac valve replacementmistake (Roberts), 930
- Central aortic pressure, noninvasive estimation of using the oscillometric method for analyzing systemic artery pulsatile blood flow: comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure
- measurements (Borow and Newburger), 879
 CHF, paroxysmal nocturnal dyspnea in: mechanisms and aminophylline therapy (Rees and Clark), 311 (Letter)
- severe, captopril therapy for: hypotensive response in presence of marked y elevated PRA (LaBarre, O'Connell, and Gunnar), 308 (Letter)
- Child, idiopathic dilation of the right atrium in a (Beder, Nihill, and McNamara), 134 (Brief comm.)
- Cholesterol, HDL-, and coronary disease, role of diet to: serum total and HDL-cholesterol in boys and men from developing and developed countries (Knuiman, West, and Hautvast), 447 (Letter)
- Chronotropy, positive, catecholamine-enhanced, without altered arrhythmogenicity: denervation supersensitivity in the canine heart (Korczyn et al.), 443 (Brief comm.)

 Circulatory collapse, cardiogenic, acute, comparative hemodynamic effects of dopamine and dobutamine in patients with (Excesse Septeman and Hodges), 202
- with (Francis, Sharma, and Hodges), 995
- Click, midsystolic, accompanying isolated bicuspid pulmonic valve (Mehlman et al.), 145 (Brief comm.)
- Clinical confirmation of ECG criteria for left atrial rhythm (Beder et al.), 848

- Coarctation of the aorta in the newborn, transluminal aortic balloon angioplasty for (Singer, Rowen, and Dorsey), 131 (Brief comm.)
- Combined intravenous administration of diazoxide and betablocking agent in acute treatment of severe hypertension or hypertensive crisis (Huysmans, Thien, and Koene), 395
- Comparative analysis of physiologic responses to three different maximal graded exercise test protocols in healthy women (Pollock et al.), 363
- heart block, long-term follow-up of young adults Complete following permanent pacemaker placement for (Bes-
- ley et al.), 332
 sequential AV pacing in right ventricular myocardial
 infarction with (Abraham and Norris), 929 (Letter)
 Complications of cardiac valve replacement: characteristic abnor-
- malities of prostheses pertaining to any or specific site (Roberts), 113
- Computed tomography as a cardiovascular diagnostic technique, the emergence of (Brundage and Lipton), 313 diagnostic role of in pericardial heart disease: congenital
 - defects, thickening, neoplasms, and effusions (Moncada et al.), 263
 - ECG synchronized, in clinical evaluation of total and regional cardiac motion: comparison of postmyocardial infarction to normal hearts by rapid sequential imaging (Masuda et al.), 230
 - in the diagnosis of pericardial cyst (Kaimal), 566 (Brief
- positron-emission, assessment of regional myocardial ischemia by (Schelbert et al.), 588
 Concealed accessory pathway, post-myocardial infarction incessant supraventricular tachycardia due to (Prystowsky et al.), 426 (Brief comm.)
- Conduction abnormalities, prognostic correlation to related hemodynamic, valvular, rhythm, and: echographicquantified severity of mitral anulus calcification (Mellino et al.), 222
- properties in right anterior accessory connections, prolonged and decremental antegrade: wide QRS antidromic tachycardia of left bundle branch block pattern without Wolff-Parkinson-White configuration in sinus
- rhythm (Gillette et al.), 66 retrograde, in supraventricular tachycardia therapy, verapamil effects on (Aryanpur-Kashani), 306 (Letter)
- Reply (Hamer, Peter, and Mandel), 306 Congenital heart diseases, other complex, and tetralogy of Fallot, the case for preoperative coronary angiography in patients with (McManus et al.), 451
 Congestive heart failure (CHF) and acute myocardial infarction,
- implications of vasodilator therapy in (Cohn), 773 in infancy: recognition and management (Artman and
 - Graham), 1040
 - paroxysmal nocturnal dyspnea in: mechanisms and ami-nophylline therapy (Rees and Clark), 311 (Letter) severe, captopril therapy for: hypotensive response in presence of markedly elevated PRA (LaBarre, O'Con-
 - nell, and Gunnar), 308 (Letter)
 - chronic, long-term hemodynamic and clinical efficacy of captopril therapy in ambulatory management of (Awan et al.), 474
 - postural hypotension during tilt with chronic captopril and diuretic therapy of (Cody, Franklin, and Laragh),
- Constriction, chronic pericardial, remote right ventricular myo-cardial infarction mimicking (Butman et al.), 912 (Brief comm.)
- Continuous electrocardiographic monitoring in patients with unstable angina pectoris: identification of high-risk subgroup with severe coronary disease, variant angina,
- and/or impaired early prognosis (Johnson et al.), 4
 Contrast left ventriculography, axial oblique, quantitative: validation of the method by demonstrating improved visualization of regional wall motion and mitral valve function with accurate volume determinations (Rogers et al.), 185
- , successful, of out-of-hospital life-threatening arrhythmias with the implanted automatic defibrilla-tor (Mirowski et al.), 147 (Brief comm.) Conversion, successful,

- Coronary angiography during acute myocardial infarction: a search for spasm (Leinbach and Gold), 768 preoperative, in patients with tetralogy of Fallot and other
 - complex congenital heart diseases, the case for (McManus et al.), 451
 - angioplasty and implications for the future, results from
 - (Gruentzig), 779 artery bypass graft, saphenous vein, angiographic documenta-tion of spontaneous spasm of (Walinsky), 290 (Brief comm.)
 - surgery, transient bundle branch block following use of hypothermic cardioplegia in: high incidence without perioperative myocardial infarction (O'Connell et al.),
 - disease, comparative effects of oral verapamil and propranolol on exercise-induced myocardial ischemia and energetics in patients with: single-blind placebo crossover evaluation using radionuclide ventriculography (Josephson et al.), 978
 heart failure secondary to, vasodilators in (Parmley, Rouleau, and Chatterjee), 625

 - left main, high predictive accuracy for detection of by
 - antilog signal processing of two-dimensional echocar-diographic images (Friedman et al.), 194
 patients and normal subjects, comparison of platelet func-tion during exercise in: potential role of platelet activation in myocardial ischemia (Mehta and Meh-
 - left anterior descending, anomalous, inadvertent division of during complete repair of tetralogy of Fallot: 22-year follow-up (Heuser, Achuff, and Brinker), 430 (Brief
 - characteristic electrocardiographic pattern indicating a critical stenosis high in in patients admitted because of impending myocardial infarction (de Zwaan, Bär, and Wellens), 730
 - pathoanatomy, right, relationship of initial and delayed right ventricular thallium-201 rest-imaging following dipyridamole-induced coronary vasodilation to (Brown et al.), 1019
 - right, rupture of in Behcet's syndrome, huge false aneurysm due to (Kaseda et al.), 569 (Brief comm.)
 - spasm: prevalence, clinical significance, and provocative testing (Conti, Feldman, and Pepine), 584 randomized double-blind comparison of nifedipine and
 - isosorbide dinitrate therapy in variant angina pectoris due to (Ginsburg et al.), 44
 - variability of electrocardiographic responses to repeated ergonovine provocation in variant angina patients with (Whittle et al.), 161
 - atherosclerosis and acute myocardial ischemic attacks, role of hypertension and downward changes of blood pres-
 - sure in the genesis of (Green), 579 bridges, isolated, clinical significance of: benign and frequent condition involving the left anterior descending artery (Kramer et al.), 283
 - bypass surgery, life-threatening exertional ventricular tachycardia in ischemic heart disease best treated by? (Cantwell), 152 (Letter)
 - calcium-a clue to angiographic underestimation of coronary luminal narrowing (Waller et al.), 1071 (Brief
 - care system, mobile, reduction of mortality from prehospital myocardial infarction by prudent patient activation of (Lewis et al.), 123
 - constriction, ergonovine, inhibition of, and variant angina pectoris, efficacy of chronic amiodarone therapy in patients with (Rutitzky, Girotti, Rosenbaum), 38
 - death, sudden, necessity of a uniform definition of: witnessed death within 1 hour of the onset of acute symptoms (Goldstein), 156
 - and myocardial metal concentrations water hardness. (Chipperfield), 1085 (Letter)
 - disease, concomitant, prevalence and significance of reversible radionuclide ischemic perfusion defects in symptomatic aortic valve disease patients with or without
 - (Pfisterer et al.), 92 mortality, recent decline in the United States: part of a general decline in mortality (Gordon), 151 (Letter)

- Coronary disease--continued
 - role of diet to HDL-cholesterol and: serum total and HDLcholesterol in boys and men from developing and developed countries (Knuiman, West, and Hautvast), 447 (Letter)
 - severe, variant angina, and/or impaired early prognosis, identification of high-risk subgroup with: continuous
 - electrocardiographic monitoring in patients with unstable angina pectoris (Johnson et al.), 4 heart disease, multiple risk functions for predicting: the con-cept, accuracy, and application (Gordon and Kannel),
 - luminal narrowing, coronary calcium—a clue to angiographic underestimation of (Waller et al.), 1071 (Brief comm.)
 - occlusion, experimental, acute, early deleterious hemorheologic changes following and salutary antihyperviscosity effect of hemodilution with stroma-free hemoglobin (Biro, Beresford-Kroeger, and Hendry), 870
 - patients, reduction of submaximal exercise myocardial oxygen demand post-walk training program in due to improved physical work efficiency (Dressendorfer et
 - with stable angina pectoris, effects of beta-adrenergic block-ade with pindolol versus placebo in (Dwyer, Pepe, and Pinkernell), 830 spasm, possible mechanisms: walk-through angina phenome
 - non demonstrated by graded exercise radionuclide ventriculography (Rizi et al.), 292 (Brief comm.)
 - rebound, possible, to abrupt nifedipine withdrawal (Kay, Biake, and Rubin), 308 (Letter) stenoses in patients, noninvasive detection and localization of:
 - comparison of resting dipyridamole and exercise thallium-201 myocardial perfusion imaging (Josephson et al.), 1008
 - vasodilation, dipyridamole-induced, initial and delayed right ventricular thallium-201 rest-imaging following: relationship to right coronary artery pathoanatomy (Brown et al.), 1019
 - vasospasm, acute allergic reaction with (Hirsh), 928 (Letter)
 - Reply (Ginsburg and Bristow), 928 role of in the pathogenesis of myocardial ischemia and angina pectoris (Gorlin), 598
- Coumarin anticoagulation, oral, long-term, prediction of hemorrhage during by excessive prothrombin ratio (Forfar), 445 (Brief comm.)
- Cultured heart cells, studies in: hypoxia, calcium fluxes, and
- inotropic state (Smith et al.), 716 Current, slow inward, potential biochemical mechanisms for regulation of: theoretical basis for drug action (Lindemann, Bailey, and Watanabe), 746
- Cyst, pericardial, computed tomography in the diagnosis of (Kaimal), 566 (Brief comm.)

D

- Death, sudden coronary, necessity of a uniform definition of: witnessed death within 1 hour of the onset of acute symptoms (Goldstein), 156
 - water hardness, and myocardial metal concentrations (Chipperfield), 1085 (Letter) high risk of, management of patients at (Lown), 689
 - in ischemic heart disease, relation of myocardial magnesium
 - deficiency to (Singh and Cameron), 449 (Letter) prevention trial, selection of an antiarrhythmic drug for a (Woosley et al.), 737
- Decline in coronary disease mortality in the United States, recent: part of a general decline in mortality (Gordon), 151 (Letter)
- Deep vein thrombosis, limb, new technique for localization and dynamic evaluation of: the 123I-fibrinogen sum-coincidence method (Ulmsten, Bernstein, and Mattson), 448 (Letter)
- Defibrillator, implanted automatic, successful conversion of outof-hospital life-threatening arrhythmias with the (Mirowski et al.), 147 (Brief comm.)
- Denervation supersensitivity in the canine heart: catecholamineenhanced positive chronotropy without altered arrhythmogenicity (Korczyn et al.), 443 (Brief

- Depolarization, myocardial, electrical-induced sustained, as a possible cause for transient ST elevation post-DC elective cardioversion (Zelinger, Falk, and Hood), 1073 (Brief comm.)
- Diabetes mellitus, increased incidence and clinical correlation of persistently abnormal technetium pyrophosphate myocardial scintigrams following acute myocardial infarction in patients with (Nicod et al.), 822
- Diagnostic promise of positron tomography (Sobel), 673 Diagnostic promise of postron tomography (Sobet), 67.

 Diastolic murmur, early, consequent apparent, shortened aortic ejection period with in mitral valve prolapse syndrome (Venetis et al.), 924 (Brief comm.)

 orthostatic blood pressure tilt-test response, increased, prognostic value of and subsequent weight gain: prospec
 - tive study of predictive factors determining borderline
- hypertensive individuals who develop sustained hypertension (Payen et al.), 379
 prolapse in the flail mitral valve syndrome: a new observation providing differentiation from the mitral valve prolapse syndrome (Cherian et al.), 1074 (Brief comm.) sound, vegetation, in mitral valve endocarditis (Valovis and Charuzi), 432 (Brief comm.)
- and beta-blocking agent, combined intravenous administration of in acute treatment of severe hypertension or hypertensive crisis (Huysmans, Thien, and Diazoxide
- Koene), 395 efficacy of slow infusion of in the treatment of severe hypertension without organ hypoperfusion (Garrett and Kaplan), 390
- Diet(s), low-sodium, will persons on become iodine deficient? (Gillum), 1084 (Letter)
 - to HDL-cholesterol and coronary disease, role of: seru: a total and HDL-cholesterol in boys and men from developing and developed countries (Knuiman, West, and Hautvast), 447 (Letter)
- Digoxin in the nonsedated chronically instrumented puppy, comparative electrophysiologic effects of (Karpawich
- vs nitroprusside, comparative hemodynamic effects of in conscious dogs with aortocaval fistula-induced chronic left ventricular volume overload and normal systolic performance (O'Rourke, Badke, and Forst), 489
- Dilation of the right atrium in a child, idiopathic (Beder, Nihill, and McNamara), 134 (Brief comm.)
- Diltiazem and combined diltiazem-digoxin in patients, intravenous, intracardiac electrophysiologic study of (Mitchell et al.), 57
- -digoxin, combined, and diltiazem in patients, intravenous, intracardiac electrophysiologic study of (Mitchell et
- Dipyridamole-induced coronary vasodilation, initial and delayed right ventricular thallium-201 rest-imaging following: relationship to right coronary artery pathoanatomy (Brown et al.), 1019
- resting, and exercise thallium-201 myocardial perfusion imaging, comparison of: noninvasive detection and localization of coronary stenoses in patients (Josephson et al.), 1008
- Dissection, aortic, in Turner's syndrome (Lie), 1077 (Brief comm.)
- Diuretic and chronic captopril therapy of severe congestive heart failure, postural hypotension during tilt with (Cody, Franklin, and Laragh), 480
- Diverse cardiovascular manifestations in Behcet's disease, recognition of the (James and Thomson), 457

 Dobutamine and dopamine, comparative hemodynamic effects of in patients with acute cardiogenic circulatory collapse (Francis, Sharma, and Hodges), 995
- Dopamine and dobutamine, comparative hemodynamic effects of in patients with acute cardiogenic circulatory collapse (Francis, Sharma, and Hodges), 995 Downward displacement of competent aortic valve prosthesis
- without left ventricular enlargement: a new sign of ascending root dissection in absence of valvular regurgitation (Gooch), 920 (Brief comm.)

 Dressler's syndrome, postmyocardial infarction (Khan), 926 (Let-
- - Reply (Kossowsky, Lyon, and Spain), 926 Second comment (Haiat), 927 Third comment (Zoneraich and Rosner), 927

- Drug action, theoretical basis for: potential biochemical mechanisms for regulation of the slow inward current (Lindemann, Bailey, and Watanabe), 746
 agents, antiarrhythmic, electrophysiologic testing of (Zipes, Prystowsky, and Heger), 610

 - antiarrhythmic, selection of an for a sudden-death-prevention
 - antiarrhytimic, selection of all for a sudden death personal rical (Woosley et al.), 737 conversion of nonsustained ventricular tachycardia to sustained ventricular tachycardia during serial electro-physiologic studies: identification of drugs that exacerbate tachycardia and potential mechanisms (Rin-
- kenberger et al.), 177 Dynamic three-dimensional echocardiographic reconstruction of the intact human left ventricle: technique and initial observations (Geiser et al.), 1056
- Dysfunction, global left ventricular, favorable hemodynamic and antiarrhythmic results in patients with: aneurysmectomy and endocardial resection for ventricular tachy-
- cardia (Martin et al.), 960

 Dysplasia, arrhythmogenic right ventricular: clinical features, diagnostic techniques, and current management (Rossi et al.), 415
- two-dimensional echocardiographic detection of (Baran et al.), 1066 (Brief comm.)

 Dyspnea, paroxysmal nocturnal, in CHF: mechanisms and ami-
- nophylline therapy (Rees and Clark), 311 (Letter)
- Dysrhythmias, ventricular, chronic, incidence and severity of after repair of tetralogy of Fallot (Kavey, Blackman, and Sondheimer), 342

- Early mobilization after uncomplicated myocardial infarction
- (West), 311 (Letter)
 ECG, noninvasive, Waller before Einthoven in the clinical development of (Besterman), 572 (Letter)
 Echocardiographic detection of circulating right atrial thrombi,
- two-dimensional (Rosenzweig and Nanda), 435 (Brief comm.)
 - two-dimensional, and diagnostic features of left ventricular papillary fibroelastoma (Ong, Nanda, and Barold), 917 (Brief comm.)
 - of arrhythmogenic right ventricular dysplasia (Baran et al.), 1066 (Brief comm.)
 - of staphylococcal vegetation attached to calcified mitral anulus (D'Cruz et al.), 295 (Brief comm.)

 - diagnosis of aortic root dissection by M-mode and two-dimensional techniques (Smuckler et al.), 897
 electrocardiographic, and angiocardiographic pre- and postoperative evaluation of primary benign intramural ventricular tumors in children (Biancaniello et al.), 852
 evaluation, two-dimensional, of LV regional wall motion abnorable to the production of analyty bidy high victoria of analyty bidy high victoria of analyty bidy high victoria.
 - malities, immediate detection of early high-risk patients with acute myocardial infarction using (Horowitz and Morganroth), 814
- features, two-dimensional, diagnostic of isolated pulmonic valve endocarditis (Mehlman et al.), 137 (Brief comm.)
 - of submitral left ventricular aneurysm (Davis et al.), 289 (Brief comm.)
- identification, two-dimensional, of intracardiac leiomyomatosis
- identification, two-dimensional, of intracardiac leiomyomatosis (Maurer and Nanda), 915 (Brief comm.)
 of sinus of Valsalva-right heart fistula due to infective endocarditis (Bardy et al.), 1068 (Brief comm.)
 images, two-dimensional, high predictive accuracy for detection of left main coronary artery disease by antilog signal processing of (Friedman et al.), 194
 reconstruction, dynamic three-dimensional, of the intact human left ventricle: technique and initial observations in patients (Geiser et al.), 1056
- tions in patients (Geiser et al.), 1056
 Echocardiography, M-mode, contrast, assessment of tricuspid regurgitation by directional analysis of right atrial systolic linear reflux echoes with (Tei, Shah, and
- Ormiston), 1025
 - diagnostic accuracy of in active infective endocarditis and prognostic implications of ultrasound-detectable veg-etations (Come, Isaacs, and Riley), 839 two-dimensional, quantification of left ventricular function by:
- consideration of factors restricting image quality (Geiser et al.), 905

- Echocardiography, two-dimensional-continued
 - utility of in the differential diagnosis of the etiology of aortic regurgitation (Imaizumi et al.), 887
- Echo-Doppler velocity and turbulence mapping, range gated, in patients with valvular aortic stenosis (Goldberg et al.), 858
- Echogram, M-mode, in differential diagnosis of VT vs SVT with aberration (Curione et al.), 1084 (Letter)
 Echographic alterations of left ventricular posterior wall contraction and relaxation patterns: preclinical abnormal segmental cardiac manifestations of thalassemia major in children on transfusion-chelation therapy (Valdes-Cruz et al.), 505
 - -determined cardiac performance, acute effects of hemodialysis on: improved contractility resulting from serum increased calcium with reduced potassium despite hypovolemic-reduced cardiac output (Chaignon et
 - end-diastolic relative wall thickness, M-mode, reliable esti-mation of peak left ventricular systolic pressure by: identification of severe valvular aortic stenosis in adult patients (Reichek and Devereux), 202
- features of posterior left ventricular wall pseudoaneurysm due to Escherichia coli endocarditis (Kessler et al.), 139 (Brief comm.)
- mitral-septal separation, detection of abnormal LV function in mitral regurgitation by (Devereux and Sachs), 1086
- Reply (Child and Perloff), 1087 -phonographic documentation of systolic honk in tricuspid prolapse (Tei, Shah, and Tanaka), 294 (Brief
- comm.) -quantified severity of mitral anulus calcification: prognostic correlation to related hemodynamic, valvular, rhythm, and conduction abnormalities (Mellino et al.), 222
- study, effects of respiration on an atrial septal aneurysm of the fossa ovale shown by (Vandenbossche and Englert), 922 (Brief comm.)
- ventricular dimensions, resting, and systolic wall stress index, relationship to: radionuclide angiographic exercise left ventricular performance in chronic aortic regurgitation (Lewis et al.), 498
- Echophonocardiographic detection, high-speed, of tricuspid valve prolapse in mitral valve prolapse with discrepancy in onset of systolic murmur (Doi et al.), 301 (Brief
- Einthoven, Waller before in the clinical development of noninvasive ECG (Besterman), 572 (Letter)
- Ejection period, aortic, shortened, with consequent apparent early diastolic murmur in mitral valve prolapse syndrome (Venetis et al.), 924 (Brief comm.)
- Electrical-induced sustained myocardial depolarization as a possible cause for transient ST elevation post-DC elective cardioversion (Zelinger, Falk, and Hood), 1073 (Brief comm.)
- Electrocardiographic (ECG) changes associated with iatrogenic left pneumothorax simulating anterior myocardial infarction (Diamond and Estes), 303 (Brief comm.)
 - ischemic, with initial nifedipine therapy of severe hyperten-
 - sion (Yagil et al.), 310 (Letter) criteria for left atrial rhythm, clinical confirmation of (Beder et al.), 848
 - echocardiographic, and angiocardiographic pre- and postoperative evaluation of primary benign intramural ventricular tumors in children (Biancaniello et al.), 852
 - monitoring, continuous, in patients with unstable angina pectoris: identification of high-risk subgroup with severe coronary disease, variant angina, and/or impaired early prognosis (Johnson et al.), 4
 - in selected unstable angina patients, vasospastic ischemic mechanism of frequent asymptomatic transient ST-T
 - changes during (Biagini et al.), 13
 systems, ambulatory, frequency response characteristics of
 and their implications for ST segment analysis
 (Bragg-Remschel, Anderson, and Winkle), 20
 - observations in severe aortic valve stenosis: correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to peak systolic transaortic pressure gradient (Siegel and Roberts), 210

- Electrocardiographic—continued
 - pattern, characteristic, indicating a critical stenosis high in left anterior descending coronary artery in patients ad-
 - mitted because of impending myocardial infarction (de Zwaan, Bär, and Wellens), 730 recordings, 24-hour Holter, accurate rapid compact analog method for quantification of frequency and duration of myocardial ischemia by semiautomated analysis of

 - of myocardial iscnemia by semiaucomated analysis of (Nademanee et al.), 802
 responses to repeated ergonovine provocation in variant angina patients with coronary artery spasm, variability of (Whittle et al.), 161
 synchronized computed tomography in clinical evaluation of total and regional cardiac motion: comparison of postmyocardial infarction to normal hearts by rapid sequential imaging (Masuda et al.), 230
- Electroencephalogram, primary ventricular fibril ation during in a psychiatric patient (Morrison et al.), 1081 (Letter)
- Electrophysiologic drug testing in prophylaxis of sporadic paroxysmal atrial fibrillation: technique, application, and efficacy in severely symptomatic preexcitation patients (Bauernfeind et al.), 941
 effects, comparative, of digoxin in the nonsedated chronically
- instrumented puppy (Karpawich et al.), 1001
- procainamide on sinus function in patients with and without sinus node disease (Goldberg et al.),
- study demonstrating triple antegrade AV nodal pathways in patients with spontaneous and/or induced supraven-
- tricular tachycardia (Swiryn et al.), 168
 intracardiac, of intravenous diltiazem and combined diltiazem-digoxin in patients (Mitchell et al.), 57
- of ventricular arrhythmias, stampede to stimulationnumerators and denominators revisited relative to (Graboys), 1089
- testing of antiarrhythmic drug agents (Zipes, Prystowsky, and Heger), 610
- Endocardial resection and aneurysmectomy for ventricular tachycardia: favorable hemodynamic and antiarrhythmic results in patients with global left ventricular dysfunction (Martin et al.), 960
- Endocarditis, Escherichia coli, echographic features of posterior left ventricular wall pseudoaneurysm due to (Kessler et al.), 139 (Brief comm.)
 - infective, active, diagnostic accuracy of M-mode echocardiography in and prognostic implications of ultrasound-detectable vegetations (Come, Isaacs, and Riley),
 - two-dimensional echocardiographic identification of sinus of Valsalva-right heart fistula due to (Bardy et al.), 1068 (Brief comm.)
 - mitral valve, vegetation diastolic sound in (Valovis and Charu-zi), 432 (Brief comm.)
 - pulmonic valve, isolated, two-dimensional echocardiographic features diagnostic of (Mehlman et al.), 137 (Brief
- Enlargement, LV, downward displacement of competent aortic valve prosthesis without: a new sign of ascending root dissection in absence of valvular regurgitation (Gooch), 920 (Brief comm.)
- Ergonovine coronary constriction, inhibition of, and variant angina pectoris, efficacy of chronic amiodarone thera-py in patients with (Rutitzky, Girotti, and Rosen-baum), 38
 - induced vasospastic angina, negative U waves and peaked T waves without ST changes during spontaneous and (Matsuguchi et al.), 918 (Brief comm.)
 - provocation, repeated, in variant angina patients with coronary artery spasm, variability of electrocardiographic responses to (Whittle et al.), 161
- Escherichia coli endocarditis, echographic features of posterior left ventricular wall pseudoaneurysm due to (Kessler et al.), 139 (Brief comm.)
- ssessment of sinoatrial node function following the Mustard operation (Hesslein et al.), 351
 - capacity, variable, in variant angina and greater exertional thallium-201 myocardial defect during vasospastic ischemic ST segment elevation than with ST depression (Shimokawa et al.), 142 (Brief comm.)

Exercise-continued

left ventricular function during, continuous noninvasive moni-toring of by thoracic impedance cardiography-auto-mated derivation of systolic time intervals (Sheps et al.), 519

performance, radionuclide angiographic, in chronic aortic regurgitation: relationship to resting echographic ventricular dimensions and systolic wall stress index

(Lewis et al.), 498

myocardial oxygen demand, submaximal, reduction of, post-walk training program in coronary patients due to improved physical work efficiency (Dressendorfer et

platelet function during in normal subjects and coronary artery disease patients, comparison of: potential role of platelet activation in myocardial ischemia (Mehta and Mehta), 49

test protocols, maximal graded, three different, comparative analysis of physiologic responses to in healthy women

(Pollock et al.), 363

Extrasystoles, ventricular, late-coupled, ventricular tachycardia initiated by: the concept of longitudinal dissociation in the microreentry pathway (Kinoshita et al.), 1090

Failure, cardiac, chronic, cardiotonic agents in the management of (Weber, Andrews, and Janicki), 639

left ventricular, hemodynamic effects of prostaglandin E1 infusion in patients with acute myocardial infarction and (Popat and Pitt), 485

Fallot, tetralogy of, and other complex congenital heart diseases, the case for preoperative coronary angiography in patients with (McManus et al.), 451

inadvertent division of an anomalous left anterior descending coronary artery during complete repair of (Heuser, Achuff, and Brinker), 430 (Brief comm.) incidence and severity of chronic ventricular dysrhythmias

after repair of (Kavey, Blackman, and Sondheimer),

Fibrillation, atrial, initiation of by spontaneous ventricular premature beats in concealed Wolff-Parkinson-White syndrome (Shen and Sung), 911 (Brief comm.)

Fibrinogen, ¹⁷³I, sum-coincidence method: new technique for localization and dynamic evaluation of limb deep vein

thrombosis (Ulmsten, Bernstein, and Mattson), 448 (Letter)

Fibroelastoma, papillary, left ventricular, two-dimensional echo-cardiographic detection and diagnostic features of (Ong, Nanda, and Barold), 917 (Brief comm.) Fistula-induced chronic left ventricular volume overload, a

caval, and normal systolic performance, comparative hemodynamic effects of digoxin vs nitroprusside in conscious dogs with (O'Rourke, Badke, and Forst), 489

right heart-sinus of Valsalva, due to infective endocarditis, two-dimensional echocardiographic identification of

(Bardy et al.), 1068 (Brief comm.)

Flail mitral valve syndrome, diastolic prolapse in the: a new observation providing differentiation from the mitral valve prolapse syndrome (Cherian et al.), 1074 (Brief comm.)

Flecainide-induced QT prolongation and ventricular tachycardia (Lui et al.), 567 (Brief comm.)

Fluorocarbons, intravenous, effects of during and without oxygen enhancement on acute myocardial ischemic injury assessed by measurement of intramyocardial gas tensions (Rude et al.), 986

Fontan graft patency by radionuclide perfusion pulmonary scan

in tricuspid atresia with previous Glenn shunt, assessment of (Covitz et al.), 1072 (Brief comm.)

Fossa ovale, atrial septal aneurysm of the, effects of respiration on shown by echographic study (Vandenbossche and Englert), 922 (Brief comm.)

Frequency response characteristics of ambulatory ECG monitoring systems and their implications for ST segment analysis (Bragg-Remschel, Anderson, and Winkle),

Friedreich's ataxia, characteristics of the cardiac hypertrophy in (Gottdiener et al.), 525

Fundamental mechanisms eliciting cardiac pain, consideration of the (Malliani and Lombardi), 575

Future implications and results from coronary angioplasty (Gruentzig), 779

Gas tensions, intramyocardial, effects of intravenous fluorocar-bons during and without oxygen enhancement on acute myocardial ischemic injury assessed by measurement of (Rude et al.), 986

GIK formulation, regular insulin dose in for acute myocardial infarction therapy (Mantle), 572 (Letter)
Glenn shunt, assessment of Fontan graft patency by radionuclide perfusion pulmonary scan in tricuspid atresia with previous (Covitz et al.), 1072 (Brief comm.)

Global left ventricular dysfunction, favorable hemodynamic and antiarrhythmic results in patients with: aneurysmectomy and endocardial resection for ventricular tachycardia (Martin et al.), 960 Glucose-insulin-potassium (GIK) formulation, regular insulin

dose in for acute myocardial infarction therapy (Man-

tle), 572 (Letter)

Graded exercise radionuclide ventriculography, walk-through angina phenomenon demonstrated by: possible coronary spasm mechanisms (Rizi et al.), 292 (Brief comm.)

test protocols, maximal, three different, comparative analysis of physiologic responses to in healthy women (Pollock et al.), 363

Graft patency, Fontan, by radionuclide perfusion pulmonary scan in tricuspid atresia with previous Glenn shunt, assessment of (Covitz et al.), 1072 (Brief comm.)

Hardness, water, myocardial metal concentrations, and sudden coronary death (Chipperfield), 1085 (Letter) HDL-cholesterol and coronary disease, role of diet to: serum total

and HDL-cholesterol in boys and men from developing and developed countries (Knuiman, West, and

Hautvast), 447 (Letter)
and serum total cholesterol in boys and men from developing
and developed countries: role of diet to HDL-cholesterol and coronary disease (Knuiman, West and Haut-

vast), 447 (Letter)

Heart as a pump, relation of myosin isozymes to the (Wikman-Coffelt, Parmley, and Mason), 934
block, complete, long-term follow-up of young adults following permanent pacemaker placement for (Besley et al.), 332

sequential AV pacing in right ventricular myocardial infarction with (Abraham and Norris), 929 (Letter)

cells, cultured, studies in: hypoxia, calcium fluxes, and inotro-pic state (Smith et al.), 716 disease(s), coronary, multiple risk functions for predicting: the concept, accuracy, and application (Gordon and Kannel), 1031

ischemic, evaluation of patients with by thallium-201 myo-cardial imaging (Pitt and Thrall), 784 newer concepts in the pathophysiology of (Kirk and Son-nenblick), 756

nuclear techniques in (Wagner), 681

potential clinical applications of sympathomimetic amines in (Goldberg and Raifer), 724

roles in clinical evaluation of: high temporal resolution synchronous volumetric scanning tomogra ay (Wood et al.), 649
relation of myocardial magnesium deficiency to sudden death in (Singh and Cameron), 449 (Letter)

other complex congenital, and tetralogy of Fallot, the case for preoperative coronary angiography in patients with (McManus et al.), 451
pericardial, diagnostic role of computed tomography in:

congenital defects, thickening, neoplasms, and effusions (Moncada et al.), 263

failure, congestive, and acute myocardial infarction, implica-tions of vasodilator therapy in (Cohn), 773 in infancy: recognition and management (Artman and

Graham), 1040

Heart failure, congestive-continued

severe, captopril therapy for: hypotensive response in presence of markedly elevated PRA (LaBarre, O'Connell, and Gunnar), 308 (Letter)

chronic, long-term hemodynamic and clinical efficacy of captopril therapy in ambulatory management of (Awan et al.), 474 postural hypotension during tilt with chronic captopril and diuretic therapy of (Cody, Franklin, and Laragh),

secondary to coronary artery disease, vasodilators in (Parmley, Rouleau, and Chatterjee), 625

Hematuria, primary renal origin of: importance of RBC casts and urinary sediment exam technique (Schifferli), 573 (Letter)

Hemodialysis, acute effects of on echographic-determined cardiac performance: improved contractility resulting from serum increased calcium with reduced potassium despite hypovolemic-reduced cardiac output (Chaignon et al.), 374

Hemodilution with stroma-free hemoglobin, early deleterious hemorheologic changes following acute experimental

coronary occlusion and salutary antihyperviscosity effect of (Biro, Beresford-Kroeger, and Hendry), 870 Hemodynamic abnormalities, prognostic correlation to related valvular, rhythm, conduction, and: echographic-quantified severity of mitral anulus calcification (Mellino et al.), 222

alterations in newborn lambs, maturational changes in cardiac muscle myosin adenosine triphosphatase activity rela-

tive to (Riemenschneider et al.), 834 effects, comparative, of dopamine and dobutamine in patients with acute cardiogenic circulatory collapse (Francis, Sharma, and Hodges), 995

of prostaglandin E, infusion in patients with acute myocardial infarction and left ventricular failure (Popat and Pitt), 485

Hemoglobin, stroma-free, early deleterious hemorheologic changes following acute experimental coronary occlusion and salutary antihyperviscosity effect of hemodilution with (Biro, Beresford-Kroeger, and Hendry),

Hemorheologic changes, early deleterious, following acute experimental coronary occlusion and salutary antihyperviscosity effect of hemodilution with stroma-free hemoglobin (Biro, Beresford-Kroeger, and Hendry), 870

therapy, beneficial, of chronic peripheral arterial disorders with pentoxifylline: results of double-blind study versus vasodilator-nylidrin (Accetto), 864

Hemorrhage during long-term oral coumarin anticoagulation by excessive prothrombin ratio, prediction of (Forfar), 445 (Brief comm.)

High risk of sudden death, management of patients at (Lown), 689

patients, early, immediate detection of with acute myocardial inferction using two-dimensional echocardiographic evaluation of left ventricular regional wall motion abnormalities (Horowitz and Morganroth), 814

High-speed echophonocardiographic detection of tricuspid valve prolapse in mitral valve prolapse with discrepancy in onset of systolic murmur (Doi et al.), 301 (Brief comm.)

Holter ECG recordings, 24-hour, accurate rapid compact analog method for quantification of frequency and duration of myocardial ischemia by semiautomated analysis of (Nademanee et al.), 802

Huge false aneurysm due to rupture of right coronary artery in Behcet's syndrome (Kaseda et al.), 569 (Brief comm.)

Hyperadrenergic-effected limb causalgia: relief by IV pharmacologic norepinephrine blockade (Hannington-Kiff), 152 (Letter)

Hypertension and downward changes of blood pressure, role of in the genesis of coronary atheroscierosis and acute myocardial ischemic attacks (Green), 579 Detection and Follow-up Program, reduced stroke incidence in

enhanced by strong support system (Langford), 1087 (Letter)

severe, efficacy of slow infusion of diazoxide in the treatment of without organ hypoperfusion (Garrett and Kaplan), Hypertension, severe-continued

ischemic ECG changes with initial nifedipine therapy of (Yagil et al.), 310 (Letter)

or hypertensive crisis, combined intravenous administration of diazoxide and beta-blocking agent in acute treat-ment of (Huysmans, Thien, and Koene), 395

sustained, prospective study of predictive factors determining borderline hypertensive individuals who develop: prognostic value of increased diastolic orthostatic blood pressure tilt-test response and subsequent weight gain (Payen et al.), 379

Hypertensive crisis or severe hypertension, combined intravenous administration of diazoxide and beta-blocking agent in acute treatment of (Huysmans, Thien, and Koene),

individuals, borderline, who develop sustained hypertension, prospective study of predictive factors determining:
prognostic value of increased diastolic orthostatic
blood pressure tilt-test response and subsequent
weight gain (Payen et al.), 379
Hypertrophic cardiomyopathy, ASH in conditions other than

(Ramsdale and Bennett), 1083 (Letter)

Hypertrophy, cardiac, in Friedreich's ataxia, characteristics of the (Gottdiener et al.), 525 Hypotension, postural, during tilt with chronic captopril and diuretic therapy of severe congestive heart failure

(Cody, Franklin, and Laragh), 480

Hypotensive response in presence of markedly elevated PRA:
captopril therapy for severe CHF (LaBarre, O'Connell, and Gunnar), 308 (Letter)
Hypothermic cardioplegia in coronary artery bypass surgery,
transient bundle branch block following use of: high

incidence without perioperative myocardial infarction

(O'Connell et al.), 85 Hypovolemic-reduced cardiac output, improved contractility resulting from serum increased calcium with reduced potassium despite: acute effects of hemodialysis on echographic-determined cardiac performance (Chaignon et al.), 374

Hypoxia, calcium fluxes, and inotropic state: studies in cultured heart cells (Smith et al.), 716

125I fibrinogen sum-coincidence method: new technique for localization and dynamic evaluation of limb deep vein thrombosis (Ulmsten, Bernstein, and Mattson), 448 (Letter)

latrogenic left pneumothorax simulating anterior myocardial infarction, ECG changes associated with (Diamond and Estes), 303 (Brief comm.)

Idiopathic dilation of the right atrium in a child (Beder, Nihill,

and McNamara), 134 (Brief comm.)
Image quality, consideration of factors restricting: quantification of left ventricular function by two-dimensional echocardiography (Geiser et al.), 905

Implanted automatic defibrillator, successful conversion of outof-hospital life-threatening arrhythmias with the (Mirowski et al.), 147 (Brief comm.)

Infancy, congestive heart failure in: recognition and management (Artman and Graham), 1040

Infarction, myocardial, acute, and congestive heart failure, implications of vasodilator therapy in (Cohn), 773

coronary angiography during: a search for spasm (Leinbach and Gold), 768 factors related to vulnerability to arrhythmias in (Gettes et

immediate detection of early high-risk patients with using two-dimensional echocardiographic evaluation of left ventricular regional wall motion abnormalities (Horowitz and Morganroth), 814

in patients with diabetes mellitus, increased incidence and clinical correlation of persistently abnormal technetium pyrophosphate myocardial scintigrams following

(Nicod et al.), 822
lethal clinical, early pronounced ST segment depression
with marked J point decline heralding: necropsyelectrocardiographic correlative study (Raunio et al.),

regular insulin dose in GIK formulation for (Mantle), 572 (Letter)

- Infarction, myocardial, acute-continued
 - role of vasodilators in the changing phases of (Swan, Shah,
 - and Rubin), 707
 - impending, characteristic electrocardiographic pattern indi-cating a critical stenosis high in left anterior descending coronary artery in patients admitted because of (de Zwaan, Bär, and Wellens), 730
 - perioperative, high incidence without: transient bundle branch block following use of hypothermic cardiople-gia in coronary artery bypass surgery (O'Connell et
 - prehospital, reduction of mortality from by prudent patient activation of mobile coronary care system (Lewis et al.), 123
 - remote right ventricular, mimicking chronic pericardial con-striction (Butman et al.), 912 (Brief comm.)
 - uncomplicated, early mobilization after (West), 311 (Let-
- Infarction, postmyocardial, Dressler's syndrome (Khan), 926 (Letter)
 - Reply (Kossowsky, Lyon, and Spain), 926 Second comment (Haiat), 927
 - Third comment (Zoneraich and Rosner), 927
- Infection, adenovirus, and Balkan nephritis (Georgescu et al.), 572 (Letter)
- Infective endocarditis, active, diagnostic accuracy of M-mode echocardiography in and prognostic implications of ultrasound-detectable vegetations (Come, Isaacs, and Riley), 839
 - two-dimensional echocardiographic identification of sinus of Valsalva-right heart fistula due to (Bardy et al.), 1068 (Brief comm.)
- Inotropic state, calcium fluxes, and hypoxia: studies in cultured heart cells (Smith et al.), 7!6
- Insulin dose in GIK formulation, regular, for acute myocardial
- infarction therapy (Mantle), 572 (Letter)
 Intracardiac electrophysiologic study of intravenous diltiazem and combined diltiazem-digoxin in patients (Mitchell et al.), 57
- leiomyomatosis, two-dimensional echocardiographic identification of (Maurer and Nanda), 915 (Brief comm.)
- Intramural ventricular tumors, primary benign, in children: pre-and postoperative ECG, echocardiographic, and angiocardiographic evaluation (Biancaniello et al.),
- Intramyocardial gas tensions, effects of intravenous fluorocarbons during and without oxygen enhancement on acute myocardial ischemic injury assessed by measurement of (Rude et al.), 986

 Intravascular laser irradiation, feasibility of for in vivo visualization and therapy of cardiocirculatory diseases (Lee et al.), 1076 (Brief comm.)
- Intravenous administration, combined, of diazoxide and betablocking agent in acute treatment of severe hypertension or hypertensive crisis (Huysmans, Thien, and Koene), 395
 - diltiazem and combined diltiazem-digoxin in patients, intracar-
 - diac electrophysiologic study of (Mitchell et al.), 57 fluorocarbons, effects of during and without oxygen enhancement on acute myocardial ischemic injury assessed by measurement of intramyocardial gas tensions (Rude
- Introduction to symposium on New strategies in management of ischemic heart disease (Levine), 583
- Iodine deficient, will persons on low-sodium diets become?

 (Gillum), 1084 (Letter)

 Irradiation, laser, intravascular, feasibility of for in vivo visualization and the apy of cardiocirculatory diseases (Lee et al.), 1076 (Brief comm.)
- Ischemia, myocardial, and angina pectoris, role of coronary vasospasm in the pathogenesis of (Gorlin), 598 by semiautomated analysis of 24-hour Holter ECG record
 - ings, accurate rapid compact analog method for quantification of frequency and duration of (Nademanee et al.), 802
 - platelet activation in, potential role of: comparison of plate-let function during exercise in normal subjects and coronary artery disease patients (Mehta and Mehta),

- Ischemia, myocardial—continued

 - regional, assessment of by positron-emission computed tomography (Schelbert et al.), 588
 role of calcium-blocking agents in treatment of cardiac arrhythmias related to (Surawicz), 698
 severe, increased ventricular fibrillation threshold with (Fisher, Sonnenblick, and Kirk), 966
- Ischemic attacks, myocardial, acute, role of hypertension and downward changes of blood pressure in genesis of coronary atherosclerosis and (Green), 579
- ECG changes with initial nifedipine therapy of severe hyper-
- tension (Yagil et al.), 310 (Letter)
 heart disease, evaluation of patients with by thallium-201
 myocardial imaging (Pitt and Thrall), 784
 life-threatening exertional ventricular tachycardia in best
 treated by coronary bypass surgery? (Cantwell), 152
 - (Letter)
 - newer concepts in the pathophysiology of (Kirk and Son-
 - nenblick), 756 nuclear techniques in (Wagner), 681 potential clinical applications of sympathomimetic amines in (Goldberg and Raifer), 724
 - roles in clinical evaluation of: high temporal resolution synchronous volumetric scanning tomography (Wood et al.), 649
- relation of myocardial magnesium deficiency to sudden death in (Singh and Cameron), 449 (Letter) mechanism, vasospastic, of frequent asymptomatic transient ST-T changes during continuous electrocardiographic monitoring in selected unstable angina patients (Biag-
- ini et al.), 13
 myocardium, kinetics of C-11 palmitic acid in acutely. Part II
 of C-11 labeled palmitic acid for the noninvasive
 evaluation of regional myocardial fatty acid metabolism with positron-computed tomography (Schön et al.), 548
- pain relief in patients with acute myocardial infarction by intravenous atenolol (Ramsdale et al.), 459 perfusion defects, radionuclide, reversible, in symptomatic
- aortic valve disease patients with or without concomitant coronary disease, prevalence and significance of (Pfisterer et al.), 92
- Isosorbide dinitrate and nifedipine therapy in variant angina pectoris due to coronary artery spasm, randomized double-blind comparison of (Ginsburg et al.), 44

 Isozymes, myosin, relation of to the heart as a pump (Wikman-Coffelt, Parmley, and Mason), 934

J point decline, marked, early pronounced ST segment depression with heralding acute lethal clinical myocardial infarction: necropsy-electrocardiographic correlative study (Raunio et al.), 32

K supplementation and Na reduction on blood pressure, independent effects of (Gillum, Prineas, and Kebede), 449 (Letter)

L

- Laser irradiation, intravascular, feasibility of for in vivo visualization and therapy of cardiocirculatory diseases (Lee et al.), 1076 (Brief comm.)
- Left anterior descending artery, benign and frequent condition involving the: clinical significance of isolated coronary bridges (Kramer et al.), 283
 - coronary artery, characteristic electrocardiographic pat-tern indicating a critical stenosis high in in patients admitted because of impending myocardial infarction (de Zwaan, Bär, and Wellens), 730
 - inadvertent division of an anomalous during complete repair of tetralogy of Fallot: 22-year follow-up (Heuser, Achuff, and Brinker), 430 (Brief comm.) atria! rhythm, clinical confirmation of ECG criteria for (Beder
- et al.), 848

186

Left-continued

bundle branch block pattern without Wolff-Parkinson-White configuration in sinus rhythm, wide QRS antidromic tachycardia of: prolonged and decremental antegrade

conduction properties in right anterior accessory con-nections (Gillette et al.), 66
main coronary artery disease, high predictive accuracy for detection of by antilog signal processing of twodimensional echocardiographic images (Friedman et

al.), 194

ventricle, intact human, dynamic three-dimensional echocardiographic reconstruction of the: technique and initial observations in patients (Geiser et al.), 1056

ventricular dysfunction, global, favorable hemodynamic and antiarrhythmic results in patients with: aneurysmectomy and endocardial resection for ventricular tachycardia (Martin et al.), 960

failure, hemodynamic effects of prostaglandin E1 infusion in patients with acute myocardial infarction and (Popat

and Pitt), 485

and Pitty, 485
function, abnormal, detection of in mitral regurgitation by
echographic mitral-septal separation (Devereux and
Sachs), 1086 (Letter)
Reply (Child and Perloff), 1087

during exercise, continuous noninvasive monitoring of by thoracic impedance cardiography-automated derivation of systolic time intervals (Sheps et al.), 519

quantification of by two-dimensional echocardiography: consideration of factors restricting image quality (Geiser et al.), 905

global and segmental function, postresuscitation, evaluation of by radionuclide ventriculography in sudden coronary death survivors of prehospital cardiac arrest: correlation to subsequent short-term prognosis (Ptacin et al.), 54

papillary fibroelastoma, two-dimensional echocardiographic detection and diagnostic features of (Ong, Nanda, and

Barold), 917 (Brief comm.)

performance, radionuclide angiographic exercise, in chronic aortic regurgitation: relationship to resting echographic ventricular dimensions and systolic wall stress index (Lewis et al.), 498

posterior wall contraction and relaxation patterns, echographic alterations of: preclinical abnormal segmental cardiac manifestations of thalassemia major in children on transfusion-chelation therapy (Valdes-Cruz et

al.), 505

systolic pressure, peak, reliable estimation of by M-mode echographic-determined end-diastolic relative wall thickness: identification of severe valvular aortic stenosis in adult patients (Reichek and Devereux), 202

volume overload, aortocaval fistula-induced chronic, and normal systolic performance, comparative hemody-namic effects of digoxin vs nitroprusside in conscious dogs with (O'Rourke, Badke, and Forst), 489

wall pseudoaneurysm, posterior, due to Escherichia coli endocarditis, echographic features of (Kessler et al.),

139 (Brief comm.)

Leiomyomatosis, intracardiac, two-dimensional echocardiographic identification of (Maurer and Nanda), 915 (Brief

Lethal arrhythmias, potentially, and stable arrhythmias, evolving

concepts of management of (Myerburg et al.), 615 Lidocaine overdose, massive (Pearl), 1083 (Letter) Life-threatening exertional ventricular tachycardia in ischemic heart disease best treated by coronary bypass surgery? (Cantwell), 152 (Letter)

Limb causalgia, hyperadrenergic-effected: relief by IV pharmacologic norepinephrine blockade (Hannington-Kiff), 152

(Letter)

Lorcainide, oral, efficacy in suppression of symptomatic ventricular tachyarrhythmias, randomized double-blind placebo controlled cross-over trial documenting (Keefe, Peters, and Winkle), 511

Low energy synchronized cardioversion of ventricular tachycar-dia and termination of ventricular fibrillation in patients using a catheter electrode: clinical transvenous cardioversion of recurrent life-threatening ventricular tachyarrhythmias (Zipes et al.), 789

Low-continued

sodium diets, will persons on become iodine deficient? (Gil-lum), 1084 (Letter)

Luminal narrowing, coronary, coronary calcium—a clue to angio-graphic underestimation of (Waller et al.), 1071 (Brief comm.)

M

M-mode and two-dimensional (echocardiographic) techniques, echocardiographic diagnosis of aortic root dissection by (Smuckler et al.), 897

echocardiography, contrast, assessment of tricuspid regurgi-tation by directional analysis of right atrial systolic linear reflux echoes with (Tei, Shah, and Ormiston),

1025

diagnostic accuracy of in active infective endocarditis and prognostic implications of ultrasound-detectable veg-etations (Come, Isaacs, and Riley), 839 am in differential diagnosis of VT vs SVT with

echogram in differential diagnosis of aberration (Curione et al.), 1084 (Letter)

echographic-determined end-diastolic relative wall thickness, reliable estimation of peak left ventricular systolic pressure by: identification of severe valvular aortic stenosis in adult patients (Reichek and Devereux).

Magnesium deficiency, myocardial, relation of to sudden death in ischemic heart disease (Singh and Cameron), 449

(Letter)

Massive lidocaine overdose (Pearl), 1083 (Letter)

Maturational changes in cardiac muscle myosin adenosine triphosphatase activity relative to hemodynamic alterations in newborn lambs (Riemenschneider et al.), 834

Maximal graded exercise test protocols, three different, in healthy women, comparative analysis of physiologic responses to (Pollock et al.), 363

Mechanisms and consequences of pulmonary edema: cardiac lung, shock lung, and principles of ventilatory therapy in adult respiratory distress syndrome (Ayres), 97 Metal concentrations, myocardial, water hardness, and sudden

coronary death (Chipperfield), 1085 (Letter) Microreentry pathway, the concept of longitudinal dissociation in

the: ventricular tachycardia initiated by late-coupled ventricular extrasystoles (Kinoshita et al.), 1090 Midsystolic click accompanying isolated bicuspid pulmonic valve

(Mehlman et al.), 145 (Brief comm.)
Mitral anular calcification, high prevalence of symptomatic bradyarrhythmias due to atrioventricular node-fascicular and sinus node-atrial disease in patients with (Nair et al.), 226

anulus calcification, echographic-quantified severity of: prog-nostic correlation to related hemodynamic, valvular, rhythm, and conduction abnormalities (Mellino et

calcified, two-dimensional echocardiographic detection of staphylococcal vegetation attached to (D'Cruz et al.), 295 (Brief comm.)

regurgitation, detection of abnormal LV function in by echographic mitral-septal separation (Devereux and Sachs), 1086 (Letter)

Reply (Child and Perloff), 1087
-septal separation, echographic, detection of abnormal LV
function in mitral regurgitation by (Devereux and Sachs), 1086 (Letter) Reply (Child and Perloff), 1087

valve endocarditis, vegetation diastolic sound in (Valovis and Charuzi), 432 (Brief comm.) function and regional wall motion with accurate volume determinations, validation of the method by demonstrating improved visualization of: quantitative axial oblique contrast left ventriculography (Rogers et al.), 185

prolapse syndrome, diastolic prolapse in the flail mitral valve syndrome—a new observation providing differentia-tion from the (Cherian et al.), 1074 (Brief comm.)

shortened aortic ejection period with consequent appar ent early diastolic murmur in (Venetis et al.), 924 (Brief comm.)

Mitral valve prolapse-continued

with discrepancy in onset of systolic murmur, high-speed echophonocardiographic detection of tricuspid valve

prolapse in (Doi et al.), 301 (Brief comm.)

prosthesis, standard Björk-Shiley unmarked tilting disc,
optimal management of suspected thrombosis of
(Olinger, Thompson, and Keelan), 440 (Brief comm.)
syndrome, flail, diastolic prolapse in the: a new observation

providing differentiation from the mitral valve pro-lapse syndrome (Cherian et al.), 1074 (Brief comm.) Mobile coronary care system, reduction of mortality from prehos-pital myocardial infarction by prudent patient activation of (Lewis et al.), 123

Mobilization, early, after uncomplicated myocardial infarction (West), 311 (Letter)

Mortality, coronary disease, recent decline in in the United States: part of a general decline in mortality (Gordon), 151 (Letter)

reduction of from prehospital myocardial infarction by prudent patient activation of mobile coronary care system (Lewis et al.), 123

Multiple risk functions for predicting coronary heart disease: the concept, accuracy, and application (Gordon and Kannel), 1031

Mustard operation, exercise assessment of sinoatrial node function following the (Hesslein et al.), 351

Myocardial depolarization, electrical-induced sustained, as a possible cause for transient ST elevation post-DC elective cardioversion (Zelinger, Falk, and Hood), 1073 (Brief comm.)

fatty acid metabolism, regional, C-11 labeled palmitic acid for noninvasive evaluation of with positron-computed tomography. I. Kinetics of C-11 palmitic acid in normal myocardium (Schön et al.), 532

II. Kinetics of C-11 palmitic acid in acutely ischemic myocardium (Schön et al.), 548

imaging, thallium-201, evaluation of patients with ischemic heart disease by (Pitt and Thrall), 784

infarction, acute, and congestive heart failure, implications of

vasodilator therapy in (Cohn), 773
and left ventricular failure, hemodynamic effects of prostaglandin E, infusion in patients with (Popat and Pitt), 485

and recurrent ventricular arrhythmias in Behcet's syn-drome (Schiff et al.), 438 (Brief comm.) coronary angiography during: a search for spasm (Lein-bach and Gold), 768

factors related to vulnerability to arrhythmias in (Gettes et

immediate detection of early high-risk patients with using two-dimensional echocardiographic evaluation of left ventricular regional wall motion abnormalities (Horowitz and Morganroth), 814

in patients with diabetes mellitus, increased incidence and clinical correlation of persistently abnormal technetium pyrophosphate myocardial scintigrams following (Nicod et al.), 822 ischemic pain relief in patients with by intravenous ateno-

lol (Ramsdale et al.), 459

lethal clinical, early pronounced ST segment depression with marked J point decline heralding: necropsyelectrocardiographic correlative study (Raunio et al.),

regular insulin dose in GIK formulation for (Mantle), 572 (Letter)

role of vasodilators in the changing phases of (Swan, Shah, and Rubin), 707

ventricular septal defect and assessment of ventricular function, first-pass radionuclide angiography in visu-alization of (Hecht and Zibelli), 436 (Brief comm.) anterior, ECG changes associated with iatrogenic left pneu-mothorax simulating (Diamond and Estes), 303

impending, characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of (de Zwaan, Bär, and Wellens), 730 perioperative, high incidence without: transient bundle

branch block following use of hypothermic cardioplegia in coronary artery bypass surgery (O'Connell et al.), 85

Myocardial infarction-continued

prehospital, reduction of mortality from by prudent patient activation of mobile coronary care system (Lewis et al.), 123

right ventricular, remote, mimicking chronic pericardial con-striction (Butman et al.), 912 (Brief comm.) sequential AV pacing in with complete heart block (Abraham and Norris), 929 (Letter)

shock, acute, successful treatment of by combined percuta-neous transluminal coronary recanalization (PTCR) and percutaneous transluminal coronary angioplasty (PTCA) (Meyer et al.), 132 (Brief comm.)

uncomplicated, early mobilization after (West), 311 (Let-

ischemia and angina pectoris, role of coronary vasospasm in the pathogenesis of (Gorlin), 598 by semiautomated analysis of 24-hour Holter ECG record-

ings, accurate rapid compact analog method for quantification of frequency and duration of (Nademanee et al.), 802

exercise-induced and energetics, comparative effects of oral verapamil and propranolol on in patients with coro-nary artery disease: single-blind placebo crossover evaluation using radionuclide ventriculography (Josephson et al.), 978

platelet activation in, potential role of: comparison of platelet function during exercise in normal subjects and coronary artery disease patients (Mehta and Mehta),

regional, assessment of by positron-emission computed tomography (Schelbert et al.), 588 role of calcium-blocking agents in treatment of cardiac arrhythmias related to (Surawicz), 698 severe, increased ventricular fibrillation threshold with

(Fisher, Sonnenblick, and Kirk), 966

ischemic attacks, acute, role of hypertension and downward changes of blood pressure in genesis of coronary atherosclerosis and (Green), 579

injury, acute, effects of intravenous fluorocarbons during and without oxygen enhancement on, assessed by mea-surement of intramyocardial gas tensions (Rude et al.), 986

magnesium deficiency to sudden death in ischemic heart disease, relation of (Singh and Cameron), 449 (Letter)

metal concentrations, water hardness, and sudden coronary death (Chipperfield), 1085 (Letter)

scintigrams, technetium pyrophosphate, persistently abnormal, increased incidence and clinical correlation of following acute myocardial infarction in patients with diabetes mellitus (Nicod et al.), 822

scintigraphy, technetium-99m-pyrophosphate, noninvasive diagnosis of cardiac amyloidosis by (Sobol et al.), 563 noninvasive (Brief comm.)

technetium-99m-pyrophosphate scintigraphy in the noninvasive diagnosis of cardiac amyloidosis, value of positive (Wizenberg et al.), 468 Myosin adenosine triphosphatase activity, cardiac muscle, rela-

tive to hemodynamic alterations in newborn lambs, maturational changes in (Riemenschneider et al.),

isozymes, relation of to the heart as a pump (Wikman-Coffelt, Parmley, and Mason), 934

N

Na reduction and K supplementation on blood pressure, independent effects of (Gillum, Prineas, and Kebede), 449 (Letter)

Necropsy-electrocardiographic correlative study of early pronounced ST segment depression with marked J point decline heralding acute lethal clinical myocardial

Negative U waves and peaked T waves without ST changes during spontaneous and ergonovine-induced vaso-spastic angina (Matsuguchi et al.), 918 (Brief comm.)

Nephritis, Balkan, and adenovirus infection (Georgescu et al), 572 (Letter)

- Nifedipine and isosorbide dinitrate therapy in variant angina pectoris due to coronary artery spasm, randomized double-blind comparison of (Ginsburg et al.), 44
 - therapy of severe hypertension, ischemic ECG changes with
- initial (Yagil et al.), 310 (Letter)
 withdrawal, possible coronary spasm rebound to abrupt (Kay,
 Blake, and Rubin), 308 (Letter)
 Nitroprusside vs digoxin, comparative hemodynamic effects of in
- conscious dogs with aortocaval fistula-induced chronic left ventricular volume overload and normal systolic performance (O'Rourke, Badke, and Forst), 489
- No cardiac catheterization before cardiac valve replacement—a mistake (Roberts), 930
- Nocturnal dyspnea, paroxysmal, in CHF: mechanisms and ami-
- nophylline therapy (Rees and Clark), 311 (Letter) Nodal block, AV, paroxysmal, symptomatic spontaneous, due to
 - localized hyperresponsiveness of the AV node to vago-tonic reflexes (Strasberg et al.), 795
 pathways, triple antegrade AV, in patients with spontaneous and/or induced supraventricular tachycardia, electrophysiologic study demonstrating (Swiryn et al.), 168
- Noninvasive detection and localization of coronary stenoses in patients: comparison of resting dipyr damole and exercise thallium-201 myocardial perfusion imaging (Josephson et al.), 1008
 - diagnosis of cardiac amyloidosis by technetium-99m-pyrophosphate myocardial scintigraphy (Sobol et al.), 563 (Brief comm.)
 - ECG, Waller before Einthoven in the clinical development of (Besterman), 572 (Letter)
 - estimation of central aortic pressure using the oscillometric method for analyzing systemic artery pulsatile blood flow: comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure measurements (Borow and Newburger), 879
 - monitoring, continuous, of left ventricular function during exercise by thoracic impedance cardiography-automated derivation of systolic time intervals (Sheps et al.), 519
- Norepinephrine blockade, relief of hyperadrenergic-effected limb causalgia by IV pharmacologic (Hannington-Kiff), 152
- Normal myocardium, kinetics of C-11 palmitic acid in. Part I of C-11 labeled palmitic acid for the noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography (Schön et al.), 532 Nuclear techniques in ischemic heart disease (Wagner), 681
- Nylidrin vasodilator, results of double-blind study versus: beneficial hemorheologic therapy of chronic peripheral arterial disorders with pentoxifylline (Accetto), 864

- Oral quinidine effects on sinus node function in sick sinus syndrome patients, assessment of (Vera, Awan, and Mason), 80
- Organ hypoperfusion, efficacy of slow infusion of diazoxide in the treatment of severe hypertension without (Garrett and Kaplan), 390
- Oscillometric method for analyzing systemic artery pulsatile blood flow, noninvasive estimation of central aortic pressure using the: comparative study of indirect systolic, diastolic and mean brachial artery pressure with simultaneous direct ascending aortic pressure measurements (Borow and Newburger), 879
- Overdose, lidocaine, massive (Pearl), 1083 (Letter) therapeutic usage, and management of complications: cardio-vascular effects of tricyclic antidepressant drugs
- (Marshall and Forker), 401 Oxygen demand, submaximal exercise myocardial, reduction of, post-walk training program in coronary patients due to improved physical work efficiency (Dressendorfer et al.), 358
 - enhancement, effects of intravenous fluorocarbons during and without on acute myocardial ischemic injury assessed measurement of intramyocardial gas tensions (Rude et al.), 986

- Pacemaker placement, permanent, for complete heart block, long-term follow-up of young adults following (Besley et al.), 332
 - therapy, chronic, in young adults, primary sick sinus syndrome as an indication for: incidence, clinical features, and long-term evaluation (Kay, Estioko, and Wiener),
- Pacing, sequential AV, in right ventricular myocardial infarction with complete heart block (Abraham and Norris), 929 (Letter)
- Pain, cardiac, consideration of the fundamental mechanisms eliciting (Malliani and Lombardi), 575
- relief, ischemic, in patients with acute myocardial infarction by intravenous atenolol (Ramsdale et al.), 459
- Palmitic acid, C-11 labeled, for noninvasive evaluation of regional myocardial fatty acid metabolism with positron-computed tomography. I. Kinetics of C-11 palmitic acid in normal myocardium (Schön et al.), 532
 - II. Kinetics of C-11 palmitic acid in acutely ischemic myocardium (Schön et al.), 548
- Papillary fibroelastoma, left ventricular, two-dimensional echo-cardiographic detection and diagnostic features of (Ong, Nanda, and Barold), 917 (Brief comm.) Paroxysmal atrial fibrillation, sporadic, electrophysiologic drug testing in prophylaxis of: technique, application, and
- efficacy in severely symptomatic patients (Bauernfeind et al.), 941 preexcitation
- AV nodal block, symptomatic spontaneous, due to localized hyperresponsiveness of the AV node to vagotonic
- reflexes (Strasberg et al.), 795 nocturnal dyspnea in CHF: mechanisms and aminophylline therapy (Rees and Clark), 311 (Letter)
- Pathophysiology, diagnosis, prevention, and treatment: current status of pulmonary thromboembolic disease (Bell and Simon), 239
- of ischemic heart disease, newer concepts in the (Kirk and Sonnenblick), 756
- Pentoxifylline, beneficial hemorheologic therapy of chronic peripheral arterial disorders with: results of double-blind study versus vasodilator-nylidrin (Accetto),
- Percutaneous transluminal coronary angioplasty (PTCA) and percutaneous transluminal coronary recanalization (PTCR), successful treatment of acute myocardial infarction shock by combined (Meyer et al.), 132 (Brief comm.)
 - recanalization (PTCR) and percutaneous transluminal coronary angioplasty (PTCA), successful treatment of acute myocardial infarction shock by combined (Meyer et al.), 132 (Brief comm.)
- Pericardial constriction, chronic, remote right ventricular myo-cardial infarction mimicking (Butman et al.), 912 (Brief comm.)
- cyst, computed tomography in the diagnosis of (Kaimal), 566 (Brief comm.)
- heart disease, diagnostic role of computed tomography in: congenital defects, thickening, neoplasms, and effu-sions (Moncada et al.), 263
- Peripheral arterial disorders, chronic, beneficial hemorheologic therapy of with pentoxifylline: results of double blind study versus vasodilator-nylidrin (Accetto), 864
- Permanent pacemaker placement for complete heart block, long-term follow-up of young adults following (Besley et al.), 332
- Pheochromocytoma, cardiac arrest in (Ouzts and Newland), 928 (Letter)
- increased 24-hour urine vanillylmandelic acid in patients without (Nanji and Biank), 1088 (Letter)
 Phonographic-echographic documentation of systolic honk in
- tricuspid prolapse (Tei, Shah, and Tanaka), 294 (Brief comm.)
- Physical work efficiency, improved, reduction of submaximal exercise myocardial oxygen demand post-walk training program in coronary patients due to (Dressendorfer et al.), 358
- Physiologic responses to three different maximal graded exercise test protocols in healthy women, comparative analysis of (Pollock et al.), 363

- Pindolol versus placebo in coronary patients with stable angina pectoris, effects of beta-adrenergic blockade with (Dwyer, Pepe, and Pinkernell), 830
- Placebo controlled crossover trial, randomized double-blind, doclacebo controlled crossover trial, randomized double-blind, doc-umenting oral lorcainide efficacy in suppression of symptomatic ventricular tachyarrhythmias (Keefe, Peters, and Winkle), 511 versus pindolol in coronary patients with stable angina pecto-ris, effects of beta-adrenergic blockade with (Dwyer, Pepe, and Pinkernell), 830
- Plasma renin activity, control, relationships to immediate blood pressure response and: prediction of sustained antihypertensive efficacy of chronic captopril therapy (Waeber et al.), 384
 - hypotensive response in presence of markedly elevated: captopril therapy for severe CHF (LaBarre, O'Con-nell, and Gunnar), 308 (Letter)
- Platelet activation in myocardial ischemia, potential role of: comparison of platelet function during exercise in normal subjects and coronary artery disease patients (Mehta and Mehta), 49
 - function during exercise in normal subjects and coronary artery disease patients, comparison of: potential role of platelet activation in myocardial ischemia (Mehta and Mehta), 49
- Pneumothorax, iatrogenic left, simulating anterior myocardial infarction, ECG changes associated with (Diamond and Estes), 303 (Brief comm.)
- Positron-computed tomography, C-11 labeled palmitic acid for the noninvasive evaluation of regional myocardial fatty acid metabolism with. I. Kinetics of C-11 palmitic acid in normal myocardium (Schön et al.), 532
 - II. Kinetics of C-11 palmitic acid in acutely ischemic myocardium (Schön et al.), 548
- -emission computed tomography, assessment of regional myo-cardial ischemia by (Schelbert et al.), 588 tomography, diagnostic promise of (Sobel), 673 Postinfarction ventricular arrhythmias—which should be treated? (Bigger, Weld, and Rolnitzky), 660
- Postmyocardial infarction, comparison of to normal hearts by rapid sequential imaging: ECG synchronized computed tomography in clinical evaluation of total and regional cardiac motion (Masuda et al.), 230
 - Dressler's syndrome (Khan), 926 (Letter) Reply (Kossowsky, Lyon, and Spain), 926 Second comment (Haiat), 927
 - Third comment (Zoneraich and Rosner), 927 incessant supraventricular tachycardia due to concealed accessory pathway (Prystowsky et al.), 426 (Brief
- comm.) Postresuscitation left ventricular global and segmental function by radionuclide ventriculography in sudden coronary death survivors of prehospital cardiac arrest, evaluation of: correlation to subsequent short-term progno-
- sis (Ptacin et al.), 54
- Postural hypotension during tilt with chronic captopril and diuretic therapy of severe congestive heart failure (Cody, Franklin, and Laragh), 480

 Post-walk training program in coronary patients, reduction of submaximal exercise myocardial oxygen demand due to improved physical work efficiency (Dressendorfer
- Potassium, reduced, with serum increased calcium despite hypovolemic-reduced cardiac output, improved contractility resulting from: acute effects of hemodialysis on echographic-determined cardiac performance (Chaignon et al.), 374
- supplementation and sodium reduction on blood pressure, independent effects of (Gillum, Prineas, and Kebede), 449 (Letter)
- Preclinical abnormal segmental cardiac manifestations of thalassemia major in children on transfusion-chelation therapy: echographic alterations of left ventricular
- posterior wall contraction and relaxation patterns
 (Valdes-Cruz et al.), 505

 Preexcitation patients, severely symptomatic, technique, application, and efficacy of electrophysiologic drug testing in prophylaxis of sporadic paroxysmal atrial fibrillation in (Bauernfeind et al.), 941

 Proposition exercises when it makes a difference? (Cabb.)
- Prehospital cardiac care: does it make a difference? (Cobb), 316

- Premature beats, ventricular, spontaneous, initiation of atrial fibrillation by in concealed Wolff-Parkinson-White syndrome (Shen and Sung), 911 (Brief comm.)
- Preoperative coronary angiography in patients with tetralogy of Fallot and other complex congenital heart diseases,
- the case for (McManus et al.), 451 blood, independent effects of Na reduction and K supplementation on (Gillum, Prineas, and Kebede), Pressure, 449 (Letter)
- gradient, peak systolic transaortic, correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to: electrocardiographic observations in severe aortic valve ste-nosis (Siegel and Roberts), 210 Prevention trial, sudden-death, selection of an antiarrhythmic
- drug for a (Woosley et al.), 737
- Primary sick sinus syndrome as an indication for chronic pacemaker therapy in young adults: incidence, clinical features, and long-term evaluation (Kay, Estioko, and Wiener), 338
 - VF during EEG in a psychiatric patient (Morrison et al.), 1081 (Letter)
- Procainamide, electrophysiologic effects of on sinus function in patients with and without sinus node disease (Goldberg et al.), 75
- Programmed stimulation, role of in assessing vulnerability to ventricular arrhythmias (Horowitz et al.), 604
- ventricular stimulation in predicting vulnerability to ventricular arrhythmias and their response to antiarrhythmic
- therapy (Mason et al.), 633
 Progress report on publication goals (Mason), 1
 Prolapse, diastolic, in the flail mitral valve syndrome: a new observation providing differentiation from the mitral valve prolapse syndrome (Cherian et al.), 1074 (Brief
 - syndrome, mitral valve, diastolic prolapse in the flail mitral valve syndrome—a new observation providing differentiation from the (Cherian et al.), 1074 (Brief
 - comm.) shortened aortic ejection period with consequent apparent early diastolic murmur in (Venetis et al.), 924 (Brief comm.)
- tricuspid, phonographic-echographic documentation of systolic honk in (Tei, Shah, and Tanaka), 294 (Brief comm.) Propranolol and verapamil, oral, comparative effects of on exer-
- cise-induced myocardial ischemia and energetics in patients with coronary artery disease: single-blind
- placebo crossover evaluation using radionuclide ven-triculography (Josephson et al.), 978

 Prostaglandin E, infusion in patients with acute myocardial infarction and left ventricular failure, hemodynamic effects of (Popat and Pitt), 485
- Prostheses (valve), characteristic abnormalities of pertaining to any or specific site: complications of cardiac valve replacement (Roberts), 113 Prothrombin ratio, excessive, prediction of hemorrhage during
- long-term oral coumarin anticoagulation by (Forfar), 445 (Brief comm.)
- Pseudoaneurysm, posterior left ventricular wall, due to Escherichia coli endocarditis, echographic features of (Kessler et al.), 139 (Brief comm.)

 Psychiatric patient, primary VF during EEG in a (Morrison et al.), 1081 (Letter)
- Publication goals, progress report on (Mason), 1 Pulmonary edema, mechanisms and consequences of: cardiac lung, shock lung, and principles of ventilatory therapy in adult respiratory distress syndrome (Ayres), 97 scan, radionuclide perfusion, assessment of Fontan graft patency by in tricuspid atresia with previous Glenn shunt (Covitz et al.), 1072 (Brief comm.)
- thromboembolic disease, current status of: pathophysiology, diagnosis, prevention, and treatment (Bell and Simon), 239
- venous connection, total anomalous: survival for 62 years without surgical intervention (McManus, Luetzeler, and Roberts), 298 (Brief comm.)
 Pulmonic valve endocarditis, isolated, two-dimensional echocardiographic features diagnostic of (Mehlman et al.), 137
- (Brief comm.)
 - isolated bicuspid, midsystolic click accompanying (Mehlman et al.), 145 (Brief comm.)

- Pulsatile blood flow, systemic artery, noninvasive estimation of central aortic pressure using the oscillometric method for analyzing: comparative study of indirect systolic, diastolic, and mean brachial artery pressure with simultaneous direct ascending aortic pressure measurements (Borow and Newburger), 879
- Pump, heart as a, relation of myosin isozymes to the (Wikman-Coffelt, Parmley, and Mason), 934

- QRS amplitude, 12-lead, relation of to peak systolic transacrtic pressure gradient, correlative necropsy study to clini-cal, hemodynamic, and ECG variables demonstrating:
- electrocardiographic observations in severe aortic valve stenosis (Siegel and Roberts), 210 antidromic tachycardia, wide, of left bundle branch block pattern without Wolff-Parkinson-White configuration in sinus rhythm: prolonged and decremental antegrade acceptation. grade conduction properties in right anterior accessory connections (Gillette et al.), 66
- QT prolongation, flecainide-induced, and ventricular tachycardia (Lui et al.), 567 (Brief comm.)
- Quinidine effects on sinus node function in sick sinus syndrome patients, assessment of oral (Vera, Awan, and Mason),

R

- Radionuclide angiocardiography, current value of for shunt quantification and management in patients with secundum
 - atrial septal defect (Hurwitz et al.), 421 angiographic exercise left ventricular performance in chronic aortic regurgitation: relationship to resting echographic ventricular dimensions and systolic wall stress index (Lewis et al.), 498
 - angiography, first-pass, in visualization of acute myocardial infarction ventricular septal defect and assessment of ventricular function (Hecht and Zibelli), 436 (Brief comm.)
 - gated biventriculograms, requential regional phase mapping of in patients with sustained ventricular tachycardia: Close correlation with electrophysiologic characteris-
 - tics (Swiryn et al.), 319 ischemic perfusion defects in symptomatic aortic valve disease patients with or without concomitant coronary disease, prevalence and significance of reversible (Pfisterer et al.), 92
 - perfusion pulmonary scan, assessment of Fontan graft patency by in tricuspid atresia with previous Glenn shunt (Covitz et al.), 1072 (Brief comm.) ventriculography, evaluation of postresuscitation left ventricu-
 - lar global and segmental function by in sudden coronary death survivors of prehospital cardiac arrest: correlation to subsequent short-term prognosis (Pta-
 - cin et al.), 54 graded exercise, walk-through angina phenomenon demonstrated by: possible coronary spasm mechanisms (Rizi et al.), 292 (Brief comm.)
 - single-blind placebo crossover evaluation of comparative effects of oral verapamil and propranolol on exercise-induced myocardial ischemia and energetics in patients with coronary artery disease using (Josephson et al.), 978
- Randomized double-blind comparison of nifedipine and isosorbide dinitrate therapy in variant angina pectoris due to coronary artery spasm (Ginsburg et al.), 44 placebo controlled crossover trial documenting oral lor
 - cainide efficacy in suppression of symptomatic ven-tricular tachyarrhythmias (Keefe, Peters, and Winkle), 511
- Range gated echo-Doppler velocity and turbulence mapping in patients with valvular aortic stenosis (Goldberg et al.),
- RBC casts and urinary sediment exam technique, importance of: primary renal origin of hematuria (Schifferli), 573 (Letter)
- Recanalization, coronary, percutaneous transluminal (PTCR) and percutaneous transluminal coronary angioplasty (PTCA), successful treatment of acute myocardial infarction shock by combined (Meyer et al.), 132 (Brief comm.)

- Reduction of mortality from prehospital myocardial infarction by prudent patient activation of mobile coronary care system (Lewis et al.), 123
- Regional myocardial ischemia, assessment of by positron-emis
 - sion computed tomography (Schelbert et al.), 588
 wall motion abnormalities, LV, immediate detection of early
 high-risk patients with acute myocardial infarction
 - using two-dimensional echocardiographic evaluation of (Horowitz and Morganroth), 814 and mitral valve function with accurate volume determinations, validation of the method by demonstrating improved visualization of: quantitative axial oblique contrast left ventriculography (Rogers et al.), 185
- Regurgitant malfunction, immediate postoperative, of Björk-Shiley aortic valve due to interfering Teflon ventricu-lar septal defect repair: correction by rotation of prosthetic anulus (Wilcox et al.), 148 (Brief comm.)
- Regurgitation, aortic, utility of two-dimensional echocardiography in the differential diagnosis of the etiology of (Imaizumi et al.), 887
 - mitral, detection of abnormal LV function in by echographic mitral-septal separation (Devereux and Sachs), 1086
 - Reply (Child and Perloff), 1087
- tricuspid, by directional analysis of right atrial systolic linear reflux echoes with contrast M-mode echocardiog-raphy, assessment of (Tei, Shah, and Ormiston),
- valvular, a new sign of ascending root dissection in absence of: downward displacement of competent aortic valve prosthesis without LV enlargement (Gooch), 920 (Brief comm.)

 Relative wall thickness, M-mode echographic-determined end-
- diastolic, reliable estimation of peak left ventricular systolic pressure by: identification of severe valvular aortic stenosis in adult patients (Reichek and Devereux), 202
- Renal origin of hematuria, primary: importance of RBC casts and urinary sediment exam technique (Schifferli), 573
- Reperfusion arrhythmias, effects of calcium channel blocking agents on (Sheehan and Epstein), 973
- Replacement, cardiac valve, no cardiac catheterization before—a
- Replacement, cardiac valve, no cardiac catheterization before—a mistake (Roberts), 930

 Respiration, effects of on an atrial septal aneurysm of the fossa ovale shown by echographic study (Vandenbossche and Englert), 922 (Brief comm.)

 Respiratory distress syndrome, adult, cardiac lung, shock lung, and principles of ventilatory therapy in: mechanisms and consequences of pulmonary edema (Ayres), 97
- Retrograde conduction in supraventricular tachycardia therapy, verapamil effects on (Aryanpur-Kashani), 306 (Let-
- Reply (Hamer, Peter, and Mandel), 306 Rhythm abnormalities, prognostic correlation to related hemody-namic, conduction, valvular, and: echographic-quantified severity of mitral anulus calcification (Mellino et
- al.), 222 left atrial, clinical confirmation of ECG criteria for (Beder et
- al.), 848
 Right atrial systolic linear reflux echoes with contrast M-mode echocardiography, assessment of tricuspid regurgita-tion by directional analysis of (Tei, Shah, and Ormiston), 1025
 - thrombi, circulating, two-dimensional echocardiographic detection of (Rosenzweig and Nanda), 435 (Brief comm.)
- atrium in a child, idiopathic dilation of the (Beder, Nihill, and McNamara), 134 (Brief comm.)
- coronary artery pathoanatomy, relationship of initial and delayed right ventricular thallium-201 rest-imaging following dipyridamole-induced coronary vasodilation
- to (Brown et al.), 1019
 rupture of in Behcet's syndrome, huge false aneurysm due
 to (Kaseda et al.), 569 (Brief comm.)
 heart fistula-sinus of Valsalva, due to infective endocarditis,
 two-dimensional echocardiographic identification of (Bardy et al.), 1068 (Brief comm.)
- ventricular dysplasia, arrhythmogenic: clinical features, diagnostic techniques, and current management (Rossi et al.), 415

- Right ventricular dysplasia, arrhythmogenic-continued two-dimensional echocardiographic detection of (Baran et al.), 1066 (Brief comm.)

 - et al.), 1066 (Brief comm.)
 myocardial infarction, remote, mimicking chronic pericardial
 constriction (Butman et al.), 912 (Brief comm.)
 sequential AV pacing in with complete heart block
 (Abraham and Norris), 929 (Letter)
 thallium-201 rest-imaging, initial and delayed, following
 dipyridamole-induced coronary vasodilation: relationship to right coronary artery pathoanatomy (Brown et
- Risk functions, multiple, for predicting coronary heart disease: the concep., accuracy, and application (Gordon and Kannel), 1031
- Rannell, 1031

 Root dissection, ascending, a new sign of in absence of valvular regurgitation: downward displacement of competent aortic valve prosthesis without left ventricular enlargement (Gooch), 920 (Brief comm.)

 Rupture of right coronary artery in Behcet's syndrome, huge false aneurysm due to (Kaseda et al.), 569 (Brief comm.)

- Saphenous vein coronary artery bypass graft, angiographic documentation of spontaneous spasm of (Walinsky), 290 (Brief comm.)
- Scanning tomography, synchronous volumetric, high temporal resolution: potential roles in clinical evaluation of ischemic heart disease (Wood et al.), 649
- Scintigraphic pattern, unique, of diffuse biventricular technetium-99m-pyrophosphate uptake in amyloid heart disease: diagnostic considerations in cardiomyopathy (Schiff et al.), 562 (Brief comm.)
 Scintigraphy, myocardial, technetium-99m-pyrophosphate, non-
- invasive diagnosis of cardiac amyloidosis by (Sobol et al.), 563 (Brief comm.)
 - positive myocardial technetium-99m-pyrophosphate, value of in the noninvasive diagnosis of cardiac amyloidosis (Wizenberg et al.), 468
- Secundum atrial septal defect, current value of radionuclide angiocardiography for shunt quantification and management in patients with (Hurwitz et al.), 421
- Sequential AV pacing in right ventricular myocardial infarction with complete heart block (Abraham and Norris), 929 (Letter)
- imaging, rapid, comparison of postmyocardial infarction to normal hearts by: ECG synchronized computed tomography in clinical evaluation of total and regional cardiac motion (Masuda et al.), 230
- regional phase mapping of radionuclide gated biventriculo-grams in patients with sustained ventricular tachycar-dia: close correlation with electrophysiologic charac-teristics (Swiryn et al.), 319
- Serial electrophysiologic studies, drug conversion of nonsustained ventricular tachycardia to sustained ventricular tachycardia during: identification of drugs that exacerbate tachycardia and potential mechanisms (Rinkenberger et al.), 177
- Serum increased calcium with reduced potassium despite hypovolemic-reduced cardiac output, improved contractili-ty resulting from: acute effects of hemodialysis on echographic-determined cardiac performance (Chaignon et al.), 374
- total and HDL-cholesterol in boys and men from developing and developed countries: role of diet to HDL-choles-terol and coronary disease (Knuiman, West, and Hautvast), 447 (Letter)
- Severe hypertension, efficacy of slow infusion of diazoxide in the treatment of without organ hypoperfusion (Garrett and Kaplan), 390
- Shock lung, cardiac lung, and principles of ventilatory therapy in adult respiratory distress syndrome: mechanisms and consequences of pulmonary edema (Ayres), 97 myocardial infarction, acute, successful treatment of by com-
- bined percutaneous transluminal coronary recanaliza-tion (PTCR) and percutaneous transluminal coronary angioplasty (PTCA) (Meyer et al.), 132 (Brief comm.)

- Shortened aortic ejection period with consequent apparent early diastolic murmur in mitral valve prolapse syndrome
- (Venetis et al.), 924 (Brief comm.)
 Shunt quantification and management in patients with secundum atrial septal defect, current value of radionuclide angiocardiography for (Hurwitz et al.), 421 Sick sinus syndrome patients, assessment of oral quinidine effects
- on sinus node function in (Vera, Awan, and Mason),
 - primary, as an indication for chronic pacemaker therapy in
- young adults: incidence, clinical features, and long-term evaluation (Kay, Estioko, and Wiener), 338 Sinoatrial node function following the Mustard operation, exer-cise assessment of (Hesslein et al.), 351
- Sinus function in patients with and without sinus node disease, electrophysiologic effects of procainamide on (Goldberg et al.), 75
 - node-atrial and atrioventricular node-fascicular disease in patients with mitral anular calcification, high prevalence of symptomatic bradyarrhythmias due to (Nair et al.), 226
 - disease, electrophysiologic effects of procainamide on sinus function in patients with and without (Goldberg et
 - function in sick sinus syndrome patients, assessment of oral quinidine effects on (Vera, Awan, and Mason), 80 Valsalva-right heart fistula due to infective endocarditis,
 - two-dimensional echocardiographic identification of
- (Bardy et al.), 1068 (Brief comm.)
 rhythm, wide QRS antidromic tachycardia of left bundle
 branch block pattern without Wolff-Parkinson-White
 configuration in: prolonged and decremental antegrade conduction properties in right anterior accessory connections (Gillette et al.), 66
- syndrome, sick, primary, as an indication for chronic pacemak-er therapy in young adults: incidence, clinical features, and long-term evaluation (Kay, Estioko, and Wiener),
- Slow inward current, potential biochemical mechanisms for regulation of: theoretical basis for drug action (Lindemann, Bailey, and Watanabe), 746
 Sodium diets, low, will persons on become iodine deficient?
 (Gillum), 1084 (Letter)
- reduction and potassium supplementation on blood pressure, independent effects of (Gillum, Prineas, and Kebede), 449 (Letter)
- Spasm, a search for: coronary angiography during acute myocar-dial infarction (Leinbach and Gold), 768
 - coronary artery: prevalence, clinical significance, and provoca-tive testing (Conti, Feldman, and Pepine), 584
 - variability of electrocardiographic responses to repeated
- ergonovine provocation in variant angina patients with (Whittle et al.), 161
 rebound, possible coronary, to abrupt nifedipine withdrawal (Kay, Blake, and Rubin), 308 (Letter)
 spontaneous, of saphenous vein coronary artery bypass graft, angiographic documentation of (Walinsky), 290 (Brief comm.)
- ST changes during spontaneous and ergonovine-induced vaso-spastic angina, negative U waves and peaked T waves without (Matsuguchi et al.), 918 (Brief comm.) elevation, transient, post-DC elective cardioversion, electrical-
- induced sustained myocardial depolarization as a pos-sible cause for (Zelinger, Falk, and Hood), 1073 (Brief comm.)
- segment analysis, frequency response characteristics of ambulatory ECG monitoring systems and the implications for (Bragg-Remschel, Anderson, and Winkle), 20
- depression, variable exercise capacity in variant angina and greater exertional thallium-201 myocardial defect during vasospastic ischemic ST segment elevation than with (Shimokawa et al.), 142 (Brief comm.) with marked J point decline, early pronounced, heralding acute lethal clinical myocardial infarction: necropsycletter active (Brief variable).
 - electrocardiographic correlative study (Raunio et al.),
- elevation, vasospastic ischemic, greater exertional thallium-201 myocardial defect during than with ST depression, variable exercise capacity in variant angina and (Shimokawa et al.), 142 (Brief comm.)

- ST-T changes, vasospastic ischemic mechanism of frequent asymptomatic transient, during continuous electrocardiographic monitoring in selected unstable angina patients (Biagini et al.), 13 Stable and potentially lethal arrhythmias, evolving concepts of
- management of (Myerburg et al.), 615
- Stampede to stimulation—numerators and denominators revisited relative to electrophysiologic study of ventricular
- arrhythmias (Graboys), 1089 Staphylococcal vegetation attached to calcified mitral anulus, two-dimensional echocardiographic detection (D'Cruz et al.), 295 (Brief comm.)
- Stenoses, coronary, in patients: noninvasive detection and localization of: comparison of resting dipyridamole and exercise thallium-201 myocardial perfusion imaging (Josephson et al.), 1008
- Stenosis, critical, high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction, characteristic electrocardiographic pattern indicating a (de Zwaan, Bär, and Wellens), 730 valvular aortic, range gated echo-Doppler velocity and turbu-
- lence mapping in patients with (Goldberg et al.), 858 Stimulation, programmed, role of in assessing vulnerability to
 - ventricular arrhythmias (Horowitz et al.), 604
 ventricular, in predicting vulnerability to ventricular
 arrhythmias and their response to antiarrhythmic
 therapy (Mason et al)., 633
- stampede to--numerators and denominators revisited relative to electrophysiologic study of ventricular arrhythmias (Graboys), 1089
- Stroke incidence, reduced in HDFP enhanced by strong support system (Langford), 1087 (Letter)
 Stroma-free hemoglobin, early deleterious hemorheologic
- changes following acute experimental coronary occlusion and salutary antihyperviscosity effect of hemodilution with (Biro, Beresford-Kroeger, and Hendry), 870
- Submaximal exercise myocardial oxygen demand, reduction of, post-walk training program in coronary patients due to improved physical work efficiency (Dressendorfer et al.), 358
- Submitral left ventricular aneurysm, two-dimensional echocar-diographic features of (Davis et al.), 289 (Brief comm.)
- Sudden coronary death, necessity of a uniform definition of: witnessed death within 1 hour of the onset of acute symptoms (Goldstein), 156
 - survivors of prehospital cardiac arrest, evaluation of post-resuscitation left ventricular global and segmental function by radionuclide ventriculography in: correlation to subsequent short-term prognosis (Ptacin et al.), 54
 - water hardness, and myocardial metal concentrations (Chipperfield), 1085 (Letter)
- death, high risk of, management of patients at (Lown), 689 in ischemic heart disease, relation of myocardial magnesium deficiency to (Singh and Cameron), 449 (Letter)
- prevention trial, selection of an antiarrhythmic drug for a (Woosley et al.), 737
- Support system, strong, reduced stroke incidence in HDFP enhanced by (Langford), 1087 (Letter)
 Supraventricular tachycardia, post-myocardial infarction incessant, due to concealed accessory pathway (Prystowsky et al.), 426 (Brief comm.)
 - spontaneous and/or induced, electrophysiologic study dem-onstrating triple antegrade AV nodal pathways in patients with (Swiryn et al.), 168
 - therapy, verapamil effects on retrograde conduction in (Aryanpur-Kashani), 306 (Letter)
- Reply (Hamer, Peter, and Mandel), 306
 vs VT with aberration, M-mode echogram in differential
 diagnosis of (Curione et al.), 1084 (Letter)
 Sympathomimetic amines: potential clinical applications in is-
- chemic heart disease (Goldberg and Raifer), 724
- Synchronous volumetric scanning tomography, high temporal resolution: potential roles in clinical evaluation of ischemic heart disease (Wood et al.), 649

 Systolic honk in tricuspid prolapse, phonographic-echographic documentation of (Tei, Shah, and Tanaka), 294 (Brief

- Systolic-continued
 - linear reflux echoes, right atrial, assessment of tricuspid regurgitation by directional analysis of with contrast M-mode echocardiography (Tei, Shah, and Ormiston),
- murmur, high-speed echophonocardiographic detection of tricuspid valve prolapse in mitral valve prolapse with discrepancy in onset of (Doi et al.), 301 (Brief comm.)
- performance, normal, comparative hemodynamic effects of digoxin vs nitroprusside in conscious dogs with aortocaval fistula-induced chronic left ventricular volume
- overload and (O'Rourke, Badke, and Forst), 489
 pressure, left ventricular, peak, reliable estimation of by
 M-mode echographic-determined end-diastolic relative wall thickness: identification of severe valvular aortic stenosis in adult patients (Reichek and Devereux), 202
- time intervals, continuous noninvasive monitoring of left ventricular function during exercise by thoracic impedance cardiography-automated derivation of (Sheps et
- transaortic pressure gradient, peak, correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to: electrocardiographic observations in severe aortic valve stenosis (Siegel and Roberts), 210
- wall stress index and resting echographic ventricular dimensions, relationship to: radionuclide angiographic exercise left ventricular performance in chronic aortic regurgitation (Lewis et al.), 498

- T waves, peaked, and negative U waves without ST changes during spontaneous and ergonovine-induced vaso-spastic angina (Matsuguchi et al.), 918 (Brief spastic
- Tachyarrhythmias, spontaneously occurring, potent suppression of versus inconsistent abolition of induced ventricular tachycardia: antiarrhythmic efficacy and electrophysiologic actions of amiodarone in patients with lifethreatening ventricular arrhythmias (Nademanee et al.), 950
 - ventricular, recurrent life-threatening, clinical transvenous cardioversion of: low energy synchronized cardioversion of ventricular tachycardia and termination of ventricular fibrillation in patients using a catheter electrode (Zipes et al.), 789
 - symptomatic, randomized double-blind placebo controlled crossover trial documenting oral lorcainide efficacy in suppression of (Keefe, Peters, and Winkle), 511
- Tachycardia, drugs that exacerbate, and potential mechanisms, identification of: drug conversion of nonsustained ventricular tachycardia to sustained ventricular tachycardia during serial electrophysiologic studies (Rinkenberger et al.), 177
 - supraventricular, post-myocardial infarction incessant, due to concealed accessory pathway (Prystowsky et al.), 426 (Brief comm.)
 - spontaneous and/or induced, electrophysiologic study dem-onstrating triple antegrade AV nodal pathways in patients with (Swiryn et al.), 168
 - therapy, supraventricular, verapamil effects on retrograde con-
 - duction in (Aryanpur-Kashani), 306 (Letter)
 Reply (Hamer, Peter, and Mandel), 306
 ventricular, and flecainide-induced QT prolongation (Lui et al.), 567 (Brief comm.)
 initiated by late-coupled ventricular extrasystoles: the con
 - cept of longitudinal dissociation in the microreentry
 - pathway (Kinoshita et al.), 1090 life-threatening exertional, in ischemic heart disease best treated by coronary bypass surgery? (Cantwell), 152 (Letter)
 - nonsustained, drug conversion to sustained ventricular tachycardia during serial electrophysiologic studies: identification of drugs that exacerbate tachycardia and potential mechanisms (Rinkenberger et al.), 177

Tachycardia, ventricular-continued

sustained, during serial electrophysiologic studies, drug conversion of nonsustained ventricular tachycardia to: identification of drugs that exacerbate tachycardia and potential mechanisms (Rinkenberger et al.),

sequential regional phase mapping of radionuclide gated biventriculograms in patients with: close correlation with electrophysiologic characteristics (Swiryn et al.),

wide QRS antidromic, of left bundle branch block pattern without Wolff-Parkinson-White configuration in sinus rhythm: prolonged and decremental antegrade conduction properties in right anterior accessory connections (Gillette et al.), 66

Technetium pyrophosphate myocardial scintigrams, persistently abnormal, increased incidence and clinical correlation

of following acute myocardial infarction in patients with diabetes mellitus (Nicod et al.), 822
-99m-pyrophosphate myocardial scintigraphy, noninvasive diagnosis of cardiac amyloidosis by (Sobol et al.), 563 (Brief comm.)

scintigraphy, positive myocardial, value of in the noninva-sive diagnosis of cardiac amyloidosis (Wizenberg et

al.), 468
uptake in amyloid heart disease, unique scintigraphic pattern of diffuse biventricular; diagnostic considerations of the control of tions in cardiomyopathy (Schiff et al.), 562 (Brief comm.)

Testing, electrophysiologic, of antiarrhythmic drug agents (Zipes, Prystowsky, and Heger), 610 provocative, prevalence, and clinical significance of coronary artery spasm (Conti, Feldman, and Pepine), 584

Tetralogy of Fallot and other complex congenital heart diseases.

the case for preoperative coronary angiography in patients with (McManus et al.), 451

inadvertent division of an anomalous left anterior descending coronary artery during complete repair of (Heuser, Achuff, and Brinker), 430 (Brief comm.) incidence and severity of chronic ventricular dysrhythmias after repair of (Kavey, Blackman, and Sondheimer),

Thalassemia major in children on transfusion-chelation therapy, preclinical abnormal segmental cardiac manifesta-tions of: echographic alterations of left ventricular posterior wall contraction and relaxation patterns (Valdes-Cruz et al.), 505

Thallium-201 myocardial defect, greater exertional, during vaso-spastic ischemic ST segment elevation than with ST

depression, variable exercise capacity in variant angina and (Shimokawa et al.), 142 (Brief comm.)

imaging, evaluation of patients with ischemic heart disease by (Pitt and Thrall), 784

perfusion imaging, exercise, and resting dipyridamole, comparison of: noninvasive detection and localization of coronary stenoses in patients (Josephson et al.),

rest-imaging, right ventricular, initial and delayed, following dipyridamole-induced coronary vasodilation: relation-ship to right coronary artery pathoanatomy (Brown et al.), 1019

Thoracic impedance cardiography-automated derivation of systolic time intervals, continuous noninvasive monitoring of left ventricular function during exercise by (Sheps et al.), 519

Three-dimensional echocardiographic reconstruction, dynamic, of the intact human left ventricle: technique and initial observations in patients (Geiser et al.), 1056

Thrombi, right atrial, circulating, two-dimensional echocardio-graphic detection of (Rosenzweig and Nanda), 435 (Brief comm.)

Thromboembolic disease, pulmonary, current status of: patho physiology, diagnosis, prevention, and treatment (Bell and Simon), 239

Thrombosis, limb deep vein, new technique for localization and dynamic evaluation of: the ¹²I-fibrinogen sum-coincidence method (Ulmsten, Bernstein, and Mattson), 448 (Letter)

of standard Björk-Shiley unmarked tilting disc mitral valve prosthesis, optimal management of suspected (Olin-ger, Thompson, and Keelan), 440 (Brief comm.)

Tilting disc mitral valve prosthesis, Björk-Shiley unmarked, standard, optimal management of suspected thrombosis of (Olinger, Thompson, and Keelan), 440 (Brief

Tilt-test response, increased diastolic orthostatic blood pressure, prognostic value of and subsequent weight gain: prospective study of predictive factors determining borderline hypertensive individuals who develop sustained hypertension (Payen et al.), 379

Tomography, computed, as a cardiovascular diagnostic technique, the emergence of (Brundage and Lipton), 313 diagnostic role of in pericardial heart disease: congenital defects, thickening, neoplasms, and effusions (Mon-cada et al.), 263

ECG synchronized, in clinical evaluation of total and regional cardiac motion: comparison of postmyocardial infarction to normal hearts by rapid sequential imaging (Masuda et al.), 230

in the diagnosis of pericardial cyst (Kaimal), 566 (Brief comm.)

positron-emission, assessment of regional myocardial ische-

mia by (Schelbert et al.), 588
positron, diagnostic promise of (Sobel), 673
synchronous volumetric scanning, high temportal resolution:
potential roles in clinical evaluation of ischemic heart
disease (Wood et al.), 649

Total anomalous pulmonary venous connection: survival for 62 years without surgical intervention (McManus, Luetzeler, and Roberts), 298 (Brief comm.)

Transacrtic pressure gradient, peak systolic, correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating relation of 12-lead QRS amplitude to: electrocardiographic observations in severe aortic valve stenosis (Siegel and Roberts), 210

Transfusion-chelation therapy, preclinical abnormal segmental cardiac manifestations of thalassemia major in chil-dren on: echographic alterations of left ventricular posterior wall contractions and relaxation patterns (Valdes-Cruz et al.), 505

Transluminal aortic balloon angioplasty for coarctation of the aorta in the newborn (Singer, Rowen, and Dorsey), 131 (Brief comm.)

atresia with previous Glenn shunt, assessment of Fontan graft patency by radionuclide perfusion pul-Tricuspid monary scan in (Covitz et al.), 1072 (Brief comm.)

prolapse, phonographic-echographic documentation of systolic honk in (Tei, Shah, and Tanaka), 294 (Brief comm.) regurgitation by directional analysis of right atrial systolic linear reflux echoes with contrast M-mode echocardiography, assessment of (Tei, Shah, and Ormiston),

valve prolapse in mitral valve prolapse with discrepancy in onset of systolic murmur, high-speed echophonocar-diographic detection of (Doi et al.), 301 (Brief comm.)

Tricyclic antidepressant drugs, cardiovascular effects of: therapeutic usage, overdose, and management of complica-tions (Marshall and Forker), 401

Tumors, ventricular, primary benign intramural, in children: pre-and postoperative ECG, echocardiographic, and angiocardiographic evaluation (Biancaniello et al.),

Turner's syndrome, aortic dissection in (Lie), 1077 (Brief comm.)

Twelve-lead QRS amplitude, relation of to peak systolic transaor-tic pressure gradient, correlative necropsy study to clinical, hemodynamic, and ECG variables demonstrating: electrocardiographic observations in severe

aortic valve stenosis (Siegel and Roberts), 210 Two-dimensional and M-mode (echocardiographic) techniques, echocardiographic diagnosis of aortic root dissection by (Smuckler et al.), 897

echocardiographic detection and diagnostic features of left ventricular papillary fibroelastoma (Ong, Nanda, and Barold), 917 (Brief comm.)

of arrhythmogenic right ventricular dysplasia (Baran et al.), 1066 (Brief comm.)

circulating right atrial thrombi (Rosenzweig and Nanda), 435 (Brief comm.)

of staphylococcal vegetation attached to calcified mitral anulus (D'Cruz et al.), 295 (Brief comm.)

Two-dimensional echocardiographic-continued

evaluation of left ventricular regional wall motion abnormalities, immediate detection of early high-risk patients with acute myocardial infarction using (Horowitz and Morganroth), 814

features diagnostic of isolated pulmonic valve endocarditis (Mehlman et al.), 137 (Brief comm.) of submitral left ventricular aneurysm (Davis et al.), 289

(Brief comm.)

identification of intracardiac leiomyomatosis (Maurer and

Nanda), 915 (Brief comm.)
of sinus of Valsalva-right heart fistula due to infective
endocarditis (Bardy et al.), 1068 (Brief comm.)
images, high predictive accuracy for detection of left main

coronary artery disease by antilog signal processing of (Friedman et al.), 194

echocardiography, quantification of left ventricular function by: consideration of factors restricting image quality (Geiser et al.), 905

utility of in the differential diagnosis of the etiology of aortic regurgitation (Imaizumi et al.), 887

U waves and peaked T waves, negative, without ST changes during spontaneous and ergonovine-induced vaso-spastic angina (Matsuguchi et al.), 918 (Brief comm.)

Ultrasound-detectable vegetations, diagnostic accuracy of M-mode echocardiography in active infective endocarditis and prognostic implications of (Come, Isaacs, and Riley), 839

Underestimation, angiographic, of coronary luminal narrowing— coronary calcium (Waller et al.), 1071 (Brief comm.)

Unstable angina patients, vasospastic ischemic mechanism of frequent asymptomatic transient ST-T changes during continuous electrocardiographic monitoring in selected (Biagini et al.), 13

pectoris: is it time to change our approach? (Rackley et al.), 154

Urinary sediment exam technique, importance of RBC casts and: primary renal origin of hematuria (Schifferli), 573

Urine vanillylmandelic acid, increased 24-hour, in patients with-out pheochromocytoma (Nanji and Blank), 1088 (Let-

- Vagotonic reflexes, symptomatic spontaneous paroxysmal AV nodal block due to localized hyperresponsiveness of the AV node to (Strasberg et al.), 795
- Valsalva, sinus of, -right heart fistula due to infective endocarditis, two-dimensional echocardiographic identification of (Bardy et al.), 1068 (Brief comm.) Valve, Björk-Shiley aortic, immediate postoperative regurgitant
- malfunction of due to interfering Teflon ventricular septal defect repair: correction by rotation of prosthetic anulus (Wilcox et al.), 148 (Brief comm.)

isolated bicuspid pulmonic, midsystolic click accompanying (Mehlman et al.), 145 (Brief comm.)

prosthesis, aortic, competent, downward displacement of without left ventricular enlargement: a new sign of ascending root dissection in absence of valvular regurgitation

(Gooch), 920 (Brief comm.) replacement, cardiac, complications of: characteristic abnormalities of prostheses pertaining to any or specific site (Roberts), 113

no cardiac catheterization before-a mistake (Roberts), 930

Valvular abnormalities, prognostic correlation to related hemodynamic, rhythm, conduction, and: echographic-quantified severity of mitral anulus calcification (Mellino et al.), 222

aortic stenosis, range gated echo-Doppler valocity and turbulence mapping in patients with (Galberg et al.), 858 severe, identification of in adult patients: reliable estimation of peak left ventricular systolic pressure by M-mode echographic-determined end-diastolic rela-

tive wall thickness (Reichek and Devereux), 202

Valvular-continued

regurgitation, a new sign of ascending root dissection in absence of: downward displacement of competent aortic valve prosthesis without LV enlargement (Gooch), 920 (Brief comm.)

Vanillylmandelic acid, urine, increased 24-hour, in patients with-out pheochromocytoma (Nanji and Blank), 1088 (Let-

ter)

Variant angina patients with coronary artery spasm, variability of electrocardiographic responses to repeated ergonovine provocation in (Whittle et al.), 161

pettoris and inhibition of ergonovine coronary constriction, efficacy of chronic amiodarone therapy in patients with (Rutitzky, Girotti, and Rosenbaum), 38 severe coronary disease, and/or impaired early prognosis, identification of high-risk subgroup with: continuous

electrocardiographic monitoring in patients with unstable angina pectoris (Johnson et al.), 4
Vasodilation, coronary, dipyridamole-induced, initial and delayed right ventricular thallium-201 rest-imaging following: relationship to right coronary artery pathoanatomy (Brown et al.), 1019
Vasodilator(s) in heart failure secondary to coronary artery disease (Parmley, Rouleau, and Chatterjee), 625
-nylidrin, results of double-blind study versus: beneficial hemotheologic therapy of chronic peripheral arterial

hemorheologic therapy of chronic peripheral arterial disorders with pentoxifylline (Accetto), 864

role of in the changing phases of acute myocardial infarction (Swan, Shah, and Rubin), 707 therapy: implications in acute myocardial infarction and con-

gestive heart failure (Cohn), 773

Vasospasm, coronary, acute allergic reaction with (Hirsh), 928 (Letter) Reply (Ginsburg and Bristow), 928

role of in the pathogenesis of myocardial ischemia and angina pectoris (Gorlin), 598

Vasospastic angina, ergonovine-induced, negative U waves and peaked T waves without ST changes during spontaneous and (Matsuguchi et al.), 918 (Brief comm.)

ischemic mechanism of frequent asymptomatic transient ST-T changes during continuous electrocardiographic moni-toring in selected unstable angina patients (Biagini et

ST segment elevation than with ST depression, variable exercise capacity in variant angina and greater exer-tional thallium-201 myocardial defect during (Shimokawa et al.), 142 (Brief comm.)

Vegetation(s) diastolic sound in mitral valve endocarditis (Valovis and Charuzi), 432 (Brief comm.)

ultrasound-detectable, diagnostic accuracy of M-mode echo-cardiography in active infective endocarditis and prognostic implications of (Come, Isaacs, and Riley),

Venous connection, pulmonary, total anomalous: survival for 62 years without surgical intervention (McManus, Luet-

zeler, and Roberts), 298 (Brief comm.) Ventilatory therapy in adult respiratory distress syndrome, cardiaclung, shock lung, and principles of: mechanisms and consequences of pulmonary edema (Ayres), 97 left, intact human, dynamic three-dimensional

Ventricle, left, echocardiographic reconstruction of the: technique and initial observations in patients (Geiser et al.), 1056

Ventricular aneurysm, submitral left, two-dimensional echocar-diographic features of (Davis et al.), 289 (Brief

arrhythmias and their response to antiarrhythmic therapy, programmed ventricular stimulation in predicting vulnerability to (Mason et al.), 633

life-threatening, antiarrhythmic efficacy and electrophysiologic actions of amiodarone in patients with: potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of induced ventric-

ular tachycardia (Nademanee et al.), 950 numerators and denominators revisisted relative to electrophysiologic study of: the stampede to stimulation (Graboys), 1089

postinfarction-which should be treated? (Bigger, Weld, and Rolnitzky), 660

recurrent, and acute myocardial infarction in Behcet's syndrome (Schiff et al.), 438 (Brief comm.)

- Ventricular arrhythmias-continued
 - role of programmed stimulation in assessing vulnerability to (Horowitz et al.), 604
- dysplasia, arrhythmogenic right: clinical features, diagnostic techniques, and current management (Rossi et al.), 415
- dysrhythmias, chronic, incidence and severity of after repair of tetralogy of Fallot (Kavey, Blackman, and Sondheimer), 342
- extrasystoles, late-coupled, ventricular tachycardia initiated by: the concept of longitudinal dissociation in the microreentry pathway (Kinoshita et al.), 1090 fibrillation, primary, during EEG in a psychiatric patient (Morrison et al.), 1081 (Letter)
- - termination of in patients using a catheter electrode, low energy synchronized cardioversion of ventricular tachycardia and: clinical transvenous cardioversion of recurrent life-threatening ventricular tachyarrhythmias (Zipes et al.), 789
- threshold, increased, with severe myocardial ischemia (Fisher, Sonnenblick, and Kirk), 966
 function, first-pass radionuclide angiography in visualization of acute myocardial infarction ventricular septal defect and assessment of (Hecht and Zibelli), 436 (Brief comm.)
- left, quantification of by two-dimensional echocardiography: consideration of factors restricting image quality (Geiser et al.), 905
- premature beats, spontaneous, initiation of atrial fibrillation by in concealed Wolff-Parkinson-White syndrome (Shen
- and Sung), 911 (Brief comm.)
 septal defect, acute myocardial infarction, and assessment of
 ventricular function, first-pass radionuclide angiography in visualization of (Hecht and Zibelli), 436 (Brief
 - repair, interfering Teflon, immediate postoperative regurgitant malfunction of Björk-Shiley aortic valve due to: correction by rotation of prosthetic anulus (Wilcox et al.), 148 (Brief comm.)
- stimulation, programmed, in predicting vulnerability to ven-tricular arrhythmias and their response to antiar-rhythmic therapy (Mason et al.), 633
- tachyarrhythmias, recurrent life-threatening, clinical transve-nous cardioversion of: low energy synchronized cardio-version of ventricular tachycardia and termination of ventricular fibrillation in patients using a catheter electrode (Zipes et al.), 789
- symptomatic, randomized double-blind placebo controlled crossover trial documenting oral lorcainide efficacy in suppression of (Keefe, Peters, and Winkle), 511 tachycardia and flecainide-induced QT prolongation (Lui et
- al.), 567 (Brief comm.)
- aneurysmectomy and endocardial resection for: favorable hemodynamic and antiarrhythmic results in patients with global left ventricular dysfunction (Martin et al.), 960
- induced, potent suppression of spontaneously occurring tachyarrhythmias versus inconsistent abolition of: antiarrhythmic efficacy and electrophysiologic actions of amiodarone in patients with life-threatening ventricular arrhythmias (Nademanee et al.), 950
- initiated by late-coupled ventricular extrasystoles: the concept of longitudinal dissociation in the microreentry pathway (Kinoshita et al.), 1090
- life-threatening exertional, in ischemic heart disease best treated by coronary bypass surgery? (Cantwell), 152 (Letter)
- low energy synchronized cardioversion of and termination of ventricular fibrillation in patients using a catheter electrode: clinical transvenous cardioversion of recurrent life-threatening ventricular tachyarrhythmias (Zipes et al.), 789

- Ventricular tachycardia—continued
 - nonsustained, drug conversion of to sustained ventricular tachycardia during serial electrophysiologic studies: identification of drugs that exacerbate tachycardia and potential mechanisms (Rinkenberger et al.), 177 sustained, sequential regional phase mapping of radionuclide
 - gated biventriculograms in patients with: close correlation with electrophysiologic characteristics (Swiryn

 - lation with electrophysiologic characteristics et al.), 319
 vs SVT with aberration, M-mode echogram in differential diagnosis of (Curione et al.), 1084 (Letter) tumors, primary benign intramural, in children: pre- and postoperative ECG, echocardiographic, and angiocardiographic evaluation (Biancaniello et al.), 852
- Ventriculography, graded exercise radionuclide, walk-through angina phenomenon demonstrated by: possible coro-nary spasm mechanisms (Rizi et al.), 292 (Brief comm.)
- left, quantitative axial oblique contrast: validation of the method by demonstrating improved visualization of regional wall motion and mitral valve function with accurate volume determinations (Rogers et al.), 185 radionuclide, evaluation of postresuscitation left ventricular global and segmental function by in sudden coronary
- death survivors of prehospital cardiac arrest: correlation to subsequent short-term prognosis (Ptacin et al.) 54
 - single-blind placebo crossover evaluation of comparative effects of oral verapamil and propranolol on exercise-induced myocardial ischemia and energetics in patients with coronary artery disease using (Joseph-
- son et al.), 978 Verapamil and propranolol, oral, comparative effects of on exer-cise-induced myocardial ischemia and energetics in patients with coronary artery disease: single-blind placebo crossover evaluation using radionuclide ventriculography (Josephson et al.), 978
 - effects on retrograde conduction in supraventricular tachycar-dia therapy (Aryanpur-Kashani), 306 (Letter)
- Reply (Hamer, Peter, and Mandel), 306 Volume overload, left ventricular, aortocaval fistula-induced chronic, and normal systolic performance, compara-tive hemodynamic effects of digoxin vs nitroprusside in conscious dogs with (O'Rourke, Badke, and Forst), 489
- Volumetric scanning tomography, synchronous, high temporal resolution: potential roles in clinical evaluation of ischemic heart disease (Wood et al.), 649

W

- Walk-through angina phenomenon demonstrated by graded exer-cise radionuclide ventriculography: possible coronary spasm mechanisms (Rizi et al.), 292 (Brief comm.) Waller before Einthoven in the clinical development of noninva-
- sive ECG (Besterman), 572 (Letter) Water hardness, myocardial metal concentrations, and sudden
- Water lardness, myocarda metal contentrations, and sudden coronary death (Chipperfield), 1085 (Letter)
 Wolff-Parkinson-White configuration in sinus rhythm, wide QRS antidromic tachycardia of left bundle branch block pattern without: prolonged and decremental antegrade conduction properties in right anterior accessor
 - ry connections (Gillette et al.), 66 syndrome, concealed, initiation of atrial fibrillation by spontaneous ventricular premature beats in (Shen and Sung), 911 (Brief comm.)

